

S-A4-5

MUS. COMP. ZOOL.
LIBRARY.

MAY 11 1966

TRANSACTIONS

HARVARD
UNIVERSITY,

OF THE

SAN DIEGO SOCIETY OF NATURAL HISTORY

VOLUME 14, No. 9, pp. 97-132

TYPE SPECIMENS OF FOSSIL INVERTEBRATES
IN THE SAN DIEGO NATURAL HISTORY MUSEUM

BY

EDWARD C. WILSON

*Curator of Marine Invertebrates
San Diego Natural History Museum*

SAN DIEGO, CALIFORNIA

PRINTED FOR THE SOCIETY

APRIL 29, 1966

MAY 1 1966

HARVARD
UNIVERSITY

TYPE SPECIMENS OF FOSSIL INVERTEBRATES
IN THE SAN DIEGO NATURAL HISTORY MUSEUM

BY

EDWARD C. WILSON

INTRODUCTION

This catalogue is published in compliance with Recommendation 72D of the International Code of Zoological Nomenclature, which states (ICZN, 1964:75): "Institutional responsibility.—Every institution in which types are deposited should . . . (4) publish lists of type-material in its possession or custody . . ."

The main part of the catalogue is arranged systematically by phyla and a few subordinate categories. Within these, the genus and species-group names, as originally published, are listed alphabetically. A simple cross-index ("see") refers the reader from one name to others if more than one name has been applied to the same type. Each entry also contains, respectively, the author of the lowest species-group name cited, a bibliographic reference for the publication in which the name appeared, the type category and type number (with literal suffixes if specimens are of more than one piece) of the specimen in the San Diego Natural History Museum, the geological age of the type and the formation from which it was obtained, and the general locality from which the type was collected. In most entries, age and formation are given as they were in the original publication.

"Missing" following a type number means that the specimen seems once to have been in the collection but it cannot be found now.

Inclusion or omission of figure references in the citation indicates that the type was or was not figured, respectively, except where this line is enclosed by parentheses. Such parentheses declare the specimen to be an unfigured plastotype and the reference applies to the specimen from which the cast was made. A question mark following a figure citation indicates uncertainty that this figure was made from the specimen in our collection.

A species index is included to facilitate location of species-group names in the catalogue. Varieties are listed by both the varietal name and the next higher species-group name.

Definitions of type categories and apposite discussions of their use in paleontology were presented by Schenk *et al.* (1956:5-9) and Bell (1962:vii-viii). Their definitions and concepts are followed here.

CATALOGUE

PROTOZOA

Amphistegina californica Cushman and Hanna

Cushman and Hanna, 1927, p. 56, pl. 6, fig. 3, 4, 5.

Syntype; 68 (fig. 3), 726 (fig. 4), 727 (fig. 5); Eocene, Rose Canyon Formation; 3.5 miles north of La Jolla, San Diego Co., Calif.

Actinocyclina aster Woodring

Woodring, 1930, p. 152, pl. 14, fig. 4, pl. 16, fig. 1, 3, pl. 17, fig. 2.

Paratype; 332 (pl. 14, fig. 4), 333 (pl. 16, fig. 1), 334 (pl. 16, fig. 3), 335 (pl. 17, fig. 2); Eocene; 332-334: Canada de los Sauces; 335: Jalama Creek, both Santa Barbara Co., Calif.

Bulimina subacuminata Cushman and R. E. Stewart

Cushman, Stewart, and Stewart, 1930, p. 65.

Paratypes; 9, 710-716; Pliocene; Bear River, Humboldt Co., Calif.

Bulimina subcalva Cushman and K. C. Stewart

Cushman, Stewart, and Stewart, 1930, p. 65.

Paratypes; 8, 705-709; Pliocene; Scotia Bluffs, Humboldt Co., Calif.

Cibicides sandiegensis Cushman and Hanna

Cushman and Hanna, 1927, p. 55, pl. 6, fig. 1?, 2?.

Syntype; 67 (fig. 1?, 2?); 725 (fig. 1?, 2?); Eocene, Rose Canyon Formation; 3.5 miles north of La Jolla, San Diego Co., Calif.

Discocyclina californica Schenk

Schenk, 1929, p. 224, pl. 28, fig. 6, pl. 29, fig. 2.

Holotype; 75 (fig. 6); paratype; 76 (fig. 2); Eocene, 1 1/4 miles northeast New Almaden Mine, Santa Clara Co., Calif.

- Discocyclina psula* Woodring
 Woodring, 1930, p. 148, pl. 14, fig. 4, pl. 17, fig. 2.
 Paratype; 332 (pl. 14, fig. 4), 335 (pl. 17, fig. 2), 336, 337, 388; Eocene; 332, 336, 337: Canada de los Sauces; 334: Jalama Creek, both Santa Barbara Co., Calif.
- Elphidium fax barbarensis* Nicol
 Nicol, 1944, p. 178.
 Paratype; 619; Pleistocene, Santa Barbara Formation; Bathhouse Beach, Santa Barbara, Santa Barbara Co., Calif.
- Elphidium hughesi* Cushman and Grant
 Cushman and Grant, 1927, p. 75.
 Paratype; 74, 732-740; Pliocene, Poncho Rico Formation; Pine Valley, Monterey Co., Calif.
- Epistomina eocenica* Cushman and Hanna
 Cushman and Hanna, 1927, p. 53, pl. 5, fig. 4, 5.
 Syntype; 63 (fig. 5), 723 (fig. 4); Eocene, Rose Canyon Formation; 3.5 miles north of La Jolla, San Diego Co., Calif.
- Eponides mexicana* (Cushman)
 Cushman and Hanna, 1927, p. 54, pl. 5, fig. 8, 9?
 Hypotype; 66 (fig. 8), 724 (fig. 9?); Eocene, Rose Canyon Formation; 3.5 miles north of La Jolla, San Diego Co., Calif.
- Gaudryina convexa* Cushman var. *sandiegensis* Cushman and Hanna
 Cushman and Hanna, 1927, p. 50, pl. 4, fig. 1.
 Holotype; 55; Eocene, Rose Canyon Formation; 3.5 miles north of La Jolla, San Diego Co., Calif.
- Gyroidina soldanii* d'Orbigny var. *octocamerata* Cushman and Hanna
 Cushman and Hanna, 1927, p. 56, pl. 5, fig. 7.
 Hypotype; 65; Eocene, Rose Canyon Formation; 3.5 miles north of La Jolla, San Diego Co., Calif.
- Marginulina* sp. ?
 Cushman and Hanna, 1927, p. 51, pl. 4, fig. 5.
 Hypotype; 59; Eocene, Rose Canyon Formation; 3.5 miles north of La Jolla, San Diego Co., Calif.
- Nodosaria (Dentalina) communis* (d'Orbigny)
 Cushman and Hanna, 1927, p. 52, pl. 4, fig. 11?, 12.
 Hypotype; 61 (fig. 12), 720 (fig. 11?); Eocene, Rose Canyon Formation; 3.5 miles north of La Jolla, San Diego Co., Calif.
- Nodosaria (Dentalina) consobrina* (d'Orbigny)
 Cushman and Hanna, 1927, p. 52, pl. 4, fig. 7, 8.
 Hypotype; 60 (fig. 7), 719 (fig. 8); Eocene, Rose Canyon Formation; 3.5 miles north of La Jolla, San Diego Co., Calif.
- Nodosaria latejugata* Gumbel
 Cushman and Hanna, 1927, p. 52, pl. 5, fig. 1-3.
 Hypotype; 62 (fig. 1), 721 (fig. 2), 722 (fig. 3); Eocene, Rose Canyon Formation; 3.5 miles north of La Jolla, San Diego Co., Calif.
- Nonion* cf. *umbilicatulus* (Montagu)
 Cushman and Hanna, 1927, p. 57, pl. 6, fig. 6.
 Hypotype; 69; Eocene, Rose Canyon Formation; 3.5 miles north of La Jolla, San Diego Co., Calif.
- Quinqueloculina triangularis* d'Orbigny
 Cushman and Hanna, 1927, p. 57, pl. 6, fig. 8?, 9?
 Hypotype; 71 (fig. 8?, 9?), 29 (fig. 8?, 9?); Eocene, Rose Canyon Formation; 3.5 miles north of La Jolla, San Diego Co., Calif.
- Robulus inornatus* d'Orbigny
 Cushman and Hanna, 1927, p. 51, pl. 4, fig. 4, 6.
 Hypotype; 58 (fig. 4), 718 (fig. 6); Eocene, Rose Canyon Formation; 3.5 miles north of La Jolla, San Diego Co., Calif.
- Robulus mexicanus* (Cushman) var. *nudicostatus* (Cushman and Hanna)
 Cushman and Hanna, 1927, p. 50, pl. 4, fig. 2?
 Hypotype; 56; Eocene, Rose Canyon Formation; 3.5 miles north of La Jolla, San Diego Co., Calif.
- Siphonina* cf. *jacksonensis* Cushman and Applin
 Cushman and Hanna, 1927, p. 53, pl. 5, fig. 6?
 Hypotype; 64; Eocene, Rose Canyon Formation; 3.5 miles north of La Jolla, San Diego Co., Calif.
- Textularia labiata* Reuss, var.
 Cushman and Hanna, 1927, p. 50, pl. 4, fig. 3.
 Hypotype; 57; Eocene, Rose Canyon Formation; 3.5 miles north of La Jolla, San Diego Co., Calif.
- Triloculina inornata* d'Orbigny
 Cushman and Hanna, 1927, p. 58, pl. 6, fig. 10, 11.
 Hypotype; 70 (fig. 10), 728 (fig. 11); Eocene, Rose Canyon Formation; 3.5 miles north of La Jolla, San Diego Co., Calif.

COELENTERATA

Dendrophylia hannibali Nomland

Nomland, 1916, p. 67.

Syntype; 926, 927; Oligocene, Lincoln Formation; 1.5 miles above Porter, Grays Harbor Co., Washington.

Oculina panzana Loel and Corey

Loel and Corey, 1932, p. 275.

Paratype; 801; Miocene, Vaqueros Formation; Carrizo Creek, west of the Carrizo ranch house, La Panza Mountains, San Luis Obispo Co., Calif.

Turbinolia clarki Quayle

Quayle, 1932, p. 100.

Paratype; 356, 357; Eocene, Domengine Formation; Parsons Peak, Fresno Co., Calif.

Turbinolia dickersoni Nomland

Quayle, 1932, p. 98, pl. 6, fig. 1-5.

Hypotype; 378 (fig. 3), 379 (fig. 1, 4), 380 (fig. 5), 381 (fig. 2); Eocene, Meganos Formation; branch of Salt Creek about 16 miles north of Coalinga, Fresno Co., Calif.

Turbinolia imbulata (Hanna)

Quayle, 1932, p. 103, pl. 6, fig. 11, 12, 15, 16.

Hypotype; 358 (fig. 15, 16), 359 (fig. 11, 12); Eocene, Rose Canyon Shales; 358: tributary to San Clemente Creek; 359: Rose Canyon, both San Diego Co., Calif.

Turbinolia pusillanima Nomland

Quayle, 1932, p. 101, pl. 6, fig. 9, 10.

Hypotype; 376; Eocene, Meganos Formation; T. 1 S., R. 1 E., Diablo Quadrangle, wash near creek below 1250-hill, Contra Costa Co., Calif.

Turbinolia sulcata Lamarck

Quayle, 1932, p. 109, pl. 6, fig. 13.

Hypotype; 382; Eocene; Parnes (Oise), France.

ANNELIDA

Serpula careyi Wiedey

Wiedey, 1928, p. 155, pl. 20, fig. 1.

Holotype; 42; Miocene, Vaqueros Formation; Junction of Cantinas Creek and Nacimiento River, San Luis Obispo Co., Calif. See: *Serpula coreyi* Wiedey.*Serpula coreyi* Wiedey

Loel and Corey, 1932, pl. 65, fig. 9.

Holotype; 42; Miocene, Vaqueros Formation; Junction of Cantinas Creek and Nacimiento River, San Luis Obispo Co., Calif. See: *Serpula careyi* Wiedey

BRACHIOPODA

Eohemithiris alexi Hertlein and Grant

Hertlein and Grant, 1944, p. 55.

Paratype; 907, 908; Eocene; near headwaters of west branch of Agua Media Creek, McKittrick Quadrangle, Temblor Range, California.

Laqueus vancouverensis dicgensis Hertlein and Grant

Hertlein and Grant, 1960, p. 97, pl. 20, fig. 16, 17.

Paratype; 349; Pliocene, San Diego Formation; Pacific Beach, San Diego, San Diego Co., Calif.

MOLLUSCA

Pelecypoda

Adontorbina cycgia Berry

Berry, 1947, p. 260 (6).

Paratype; 320, 321; Pleistocene; "Hilltop Quarry", San Pedro, Los Angeles Co., Calif.

Amiantis calloza (Conrad)

Grant and Gale, 1931, p. 348, pl. 17, fig. 7, 9, 12, 13.

Hypotype; 162 (fig. 7), 163 (fig. 9), 165 (fig. 12), 166 (fig. 13); Pliocene, Pico Formation; 162: Holser Canyon; 163: west of Fernando Pass; 165-166: Elsmere Canyon, all Los Angeles Co., Calif.

Amiantis callosa (Conrad) variety *stalderi* (Clark)

Grant and Gale, 1931, p. 349, pl. 17, fig. 8a, 8b.

Hypotype; 164; Miocene; well core near Bakersfield, Kern Co., Calif.

Anatina (Raeta) plicatella (Lamarck) variety *longior* Grant and Gale

Grant and Gale, 1931, p. 408, pl. 23, fig. 1a, 1b.

Holotype; 192; Miocene; 4525 feet deep from oil well northwest of Bakersfield, Kern Co., Calif.

Anomia peruviana d'Orbigny

Grant and Gale, 1931, p. 240, pl. 12, fig. 2.

Hypotype; 118; Pleistocene, edge of mesa west of Newport, Los Angeles Co., Calif.

- Anomia vaquerosensis* Loel and Corey
 Loel and Corey, 1932, p. 203.
 Paratype; 782; Miocene, Vaqueros Formation; along NE-SW ridge west of mouth of Wiley Canyon, Ventura Co., Calif.
- Antigona carizzoensis* Loel and Corey
 Loel and Corey, 1932, p. 221.
 Paratype; 784-786; Miocene, Vaqueros Formation; "V" bend in Carrizo Creek, San Luis Obispo Co., Calif.
- Antigona vaquerosensis* Loel and Corey
 Loel and Corey, 1932, p. 221.
 Paratype; 787; Miocene, Vaqueros Formation; divide between Reliz and Vaqueros canyons, Monterey Co., Calif.
- Arca (Andara) santaclarana* Loel and Corey
 Loel and Corey, 1932, p. 184.
 Paratype; 776; Miocene, Vaqueros Formation; along NE-SW ridge west of mouth of Wiley Canyon, Ventura Co., Calif.
- Arca (Andara) santana* Loel and Corey
 Loel and Corey, 1932, p. 185.
 Paratype; 777; Miocene, Vaqueros Formation; west side of Plano Trabuco opposite south end of hill on plain, Santa Ana Mountains, Orange Co., Calif.
- Arca (Arca) multicostata* Sowerby variety *camiloensis* Osmont
 Grant and Gale, 1931, p. 139, fig. 5a-c.
 Hypotype; 85; Pliocene, Pico Formation; southeast of Pico Canyon, Los Angeles Co., Calif.
- Arca (Barbatia) strongi* Loel and Corey
 Loel and Corey, 1932, p. 183.
 Paratype; 775; Miocene, Vaqueros Formation; west side of Laguna Canyon on spur in large turn, Orange Co., Calif.
- Arca galei* Wiedey
 Wiedey, 1928, p. 129, pl. 13, fig. 8.
 Holotype; 21; Miocene; Temblor Formation; Benedict Canyon, Santa Monica Mtns., Los Angeles Co., Calif.
- Arca hamelini* Wiedey
 Wiedey, 1928, p. 126, pl. 13, fig. 2.
 Holotype; 18; Miocene, Vaqueros Formation; Little Sespe Creek, Ventura Co., Calif.
- Arca impavida* Wiedey
 (Wiedey, 1928, p. 130, pl. 14, fig. 2, 3.)
 Plastoholotype; 368, 369; Miocene, Temblor Formation; Barker's Ranch, Kern Co., Calif.
- Arca lakei* Wiedey
 Wiedey, 1928, p. 127, pl. 13, fig. 4, 5.
 Holotype; 19; Miocene, Temblor Formation; 2 miles west of Simmler, San Luis Obispo Co., Calif.
- Arca (Navicula) terminumbonis* Grant and Gale
 Grant and Gale, 1931, p. 142, pl. 1, fig. 18a-c, 19a-c, 20a-c, 22.
 Holotype; 79 (fig. 18a-c), paratype; 80 (fig. 19a-c), 81 (fig. 20a-c), 82 (fig. 22), 83, 741; Pliocene, Pico Formation; Elsmere Canyon, Los Angeles Co., Calif.
- Arca perdiparis* Wiedey
 Wiedey, 1928, p. 131, pl. 13, fig. 6.
 Holotype; 23; Miocene, Monterey Formation; Santa Cruz Mountains, Monterey Co., Calif.
- Arca procumbens* Wiedey
 Wiedey, 1928, p. 132, pl. 13, fig. 10.
 Syntype; 24; Miocene; 5 miles north of Yaquina Head, Lincoln Co., Oregon.
- Arca rivularia* Wiedey
 Wiedey, 1928, p. 128, pl. 13, fig. 3.
 Holotype; 20; Miocene, Temblor Formation; 2 miles west of Simmler, San Luis Obispo Co., Calif.
- Arca sespecensis* Wiedey
 Wiedey, 1928, p. 125, pl. 13, fig. 1.
 Holotype; 17; Miocene, Vaqueros Formation; Little Sespe Creek, Ventura Co., Calif.
- Arca vancouverensis* Meek
 (Meek, 1857, p. 40.)
 Plastoparatype; 624; Cretaceous, Komoks Formation; Vancouver Island, British Columbia, Canada.
 See: *Parallelodon (Nanowanis) vancouverensis* (Meek)
- Atrina stephani* Hanna
 Hanna, 1926, p. 461, pl. 27, fig. 3, 4.
 Holotype; 2; Pliocene, Imperial Formation; Coyote Mountain, Imperial Co., Calif.

Basterotia (Basterotella) hertleinii Durham

Emerson and Herlein, 1964, p. 355, fig. 4g-4j.

Hypotype; 811 (4g, 4h), 812 (4i), 813 (4j); Pliocene; Puerto Ballandra, Isla Carmen, Baja California, Mexico.

Cardita (Carditamera) carpenteri Lamy

Lamy, 1921, p. 264.

See: *Glans minuscula* Grant and Gale*Cardita ventricosa* Gould

Grant and Gale, 1931, p. 272, pl. 13, fig. 9a, 9b, 11.

Hypotype; 124 (fig. 11), 125 (fig. 9a, 9b); 124: Pliocene, Pico Formation; Sulphur Canyon, Ventura Co., Calif.; 125: Pleistocene, ridge between Casitas Creek and the sea, Ventura Co., Calif.

Cardium arcumbona Wiedey

Wiedey, 1928, p. 142, pl. 17, fig. 5.

Holotype; 31; Miocene, Temblor Formation; 2 miles west of Simmler, San Luis Obispo Co., Calif.

Cardium schencki Wiedey

Wiedey, 1928, p. 143, pl. 17, fig. 3, 4.

Paratype; 32; Miocene, Temblor Formation; 2 miles south of Calabasas, Los Angeles Co., Calif.

Chione juanensis Loel and Corey

Loel and Corey, 1932, p. 223.

Paratype; 788; Miocene, Vaqueros Formation; near head of west branch of Anderson Creek, San Luis Obispo Co., Calif.

Chione sechurana Pilsbry and Olsson

Pilsbry and Olsson, 1935, p. 17.

Paratype; 580, 928; Pleistocene; Tric Trac Point, near Bayover, Bay of Sechura, Peru.

Chione valentinei Wiedey

Wiedey, 1929, p. 282, pl. 32, fig. 1.

Paratype; 53; Miocene, Temblor Formation (?); 2 miles south of Mayfield, Santa Clara Co., Calif.

Chlamys (Argopecten) abietis (E. K. Jordan and Hertlein)

Emerson and Herlein, 1964, p. 354, fig. 4a-4e.

Hypotype; 805 (4a), 806 (4b), 807 (4c), 808 (4d), 809 (4e); 805, 808, 809: Pliocene, Isla San José; 806, 807: Pliocene; Isla Monserrate, both Baja California, Mexico.

Clementia (?) elongata Wiedey

Wiedey, 1928, p. 147, pl. 18, fig. 6.

Holotype; 35; Miocene; Temblor Formation; 2 miles west of Simmler, San Luis Obispo Co., Calif.

Clementia inequalis Wiedey

Wiedey, 1928, p. 146, pl. 18, fig. 4.

Holotype; 34; Miocene, Vaqueros Formation; South Mountain, Ventura Co., Calif.

Clementia (Compsoomya) subdiaphana Carpenter

Grant and Gale, 1931, p. 334, pl. 17, fig. 10a, 10b.

Hypotype; 167; Pleistocene, San Pedro Formation; Deadman Island, Los Angeles Co., Calif.

Corbula (Corbula) gibbiformis Grant and Gale

Grant and Gale, 1931, p. 420, pl. 19, fig. 4-6.

Holotype; 172 (fig. 5); paratype; 171 (fig. 4), 173 (fig. 6); Pliocene, Etchegoin Formation; 3951-52 feet deep in oil well, Kern Co., Calif.

Corbula (Lentidium) luteola Carpenter

Grant and Gale, 1931, p. 421, pl. 19, fig. 2, 7.

Hypotype; 168 (fig. 2), 169 (fig. 7); Pleistocene; south of Seal Cliff Station, Ventura Co., Calif.

Crassatellites antillarum (Reeves)

Grant and Gale, 1931, p. 271, pl. 13, fig. 7a, 7b.

Hypotype; 122; Pliocene; near Santa Rosalia, Baja California, Mexico.

Cryptomya californica (Conrad)

Grant and Gale, 1931, p. 417, pl. 21, fig. 7, 11.

Hypotype; 185 (fig. 7), 315 (fig. 11); Pleistocene; 185: southwest of Goleta, Santa Barbara Co., Calif.; 315: Barlow Canyon, Ventura Co., Calif.

Cumingia lamellosa Sowerby

Grant and Gale, 1931, p. 378, pl. 14, fig. 23, pl. 19, fig. 1.

Hypotype; 143; Pliocene; Maria Ygnacia Creek, Santa Barbara Co., Calif.

Donax gouldii Dall

Grant and Gale, 1931, p. 380, pl. 13, fig. 12.

Hypotype; 127; Pleistocene; west of Newport, Los Angeles Co., Calif.

Dosinia (Dosimidia) margaritana (Wiedey) *projecta* Loel and Corey

Loel and Corey, 1932, p. 217.

Paratype; 783; Miocene, Vaqueros Formation; head of north fork of Corral de Piedra Creek, about 5 miles east of San Luis Obispo, San Luis Obispo Co., Calif.

- Dosinia margaritana* Wiedey
 Wiedey, 1928, p. 145, pl. 18, fig. 1, 3.
 Holotype; 33; Miocene, Vaqueros Formation; 0.4 miles east of La Panza, San Luis Obispo Co., Calif.
- Dosinia ponderosa* (Gray) variety *jacalitosana* Arnold
 Grant and Gale, 1931, p. 352, pl. 15, fig. 2a, 2b, 3.
 Hypotype; 144 (fig. 2a, 2b), 145 (fig. 3); Pliocene, San Diego Formation; Balboa Park, San Diego, San Diego Co., Calif.
- Dosinia ponderosa* (Gray) variety *longidens* Grant and Gale
 Grant and Gale, 1931, p. 353, pl. 15, fig. 4.
 Holotype; 146; Miocene; core from oilwell northwest of Bakersfield, Kern Co., Calif.
- Glans minuscula* Grant and Gale
 Grant and Gale, 1931, p. 277, pl. 13, fig. 10a, 10b.
 Holotype; 126; Pleistocene; Seal Cliff, Ventura Co., Calif.
- See: *Cardita (Carditamera) carpenteri* Lamy
- Glycymeris septentrionalis* (Middendorff)
 Grant and Gale, 1931, p. 134, pl. 1, fig. 21a-b.
 Hypotype; 84; Pleistocene; ridge at northwest corner El Conejo land grant, Ventura Co., Calif.
- Irus lamellifer* (Conrad) variety *prelamellifer* Grant and Gale
 Grant and Gale, 1931, p. 332, pl. 18, fig. 7.
 Holotype; 157; Miocene; well core near Bakersfield, Kern Co., Calif.
- Lacvicardium (Cerastoderma) corbis* (Martyn)
 Grant and Gale, 1931, p. 307, pl. 19, fig. 17.
 Hypotype; 176; Recent; Seven Mile Beach, San Mateo Co., Calif.
- Lacvicardium (Nemocardium) centifolium* (Carpenter)
 Grant and Gale, 1931, p. 311, pl. 19, fig. 9, 10.
 Hypotype; 174 (fig. 9), 175 (fig. 10); Pliocene, Etchegoin Formation; 3951-52 feet deep in oil well, Kern Co., Calif.
- Lucina (Here) excavata* Carpenter
 Grant and Gale, 1931, p. 290, pl. 14, fig. 2, 5, 10.
 Hypotype; 131 (fig. 2), 132 (fig. 3), 133 (fig. 10); Pliocene, Pico Formation; Holser Canyon, Los Angeles Co., Calif.
- Lucina (Miltha) xantusi* (Dall)
 Grant and Gale, 1931, p. 291, pl. 14, fig. 20a, 20b.
 Hypotype; 140a (fig. 20a, 20b), 140b; Pliocene, Pico Formation; Holser Canyon, Los Angeles Co., Calif.
- Lucina (Myrtea) acutilineata* Conrad
 Grant and Gale, 1931, p. 286, pl. 14, fig. 22a, 22b.
 Hypotype; 141; Pleistocene; San Pedro, Los Angeles Co., Calif.
- Lucina (Myrtea) californica* Conrad
 Grant and Gale, 1931, p. 285, pl. 14, fig. 15a, 15b, 21a, 21b.
 Hypotype; 138 (fig. 15a, 15b), 142 (fig. 21a, 21b); 138: Pleistocene, San Pedro, Los Angeles Co., Calif.; 142: Pliocene, Santa Barbara, Santa Barbara Co., Calif.
- Lucina (Myrtea) mutallii* Conrad
 Grant and Gale, 1931, p. 288, pl. 14, fig. 18.
 Hypotype; 139; Pliocene, Pico Formation; east of Fernando Pass, Los Angeles Co., Calif.
- Lucina (Myrtea) tenuisculpta* Carpenter variety *approximata* (Dall)
 Grant and Gale, 1931, p. 289, pl. 14, fig. 8a, 8b.
 Hypotype; 136; Pleistocene, San Pedro Formation; San Pedro, Los Angeles Co., Calif.
- Macoma balthica* (Linnaeus)
 Grant and Gale, 1931, p. 371, pl. 14, fig. 6a, 6b, pl. 20, fig. 7a, 7b.
 Hypotype; 134a (fig. 6a, 7a), 134b (fig. 6b, 7b); Pleistocene; southwest of Goleta, Santa Barbara Co., Calif.
- Macoma copelandi* Wiedey
 Wiedey, 1928, pl. 149, pl. 19, fig. 2.
 Holotype; 37; Miocene, Temblor Formation; 2 miles west of Simmler, San Luis Obispo Co., Calif.
- Macoma moesta* (Deshayes) variety *aeolasta* Dall
 Grant and Gale, 1931, p. 371, pl. 14, fig. 7.
 Hypotype; 135; Pleistocene; near Goleta, Santa Barbara Co., Calif.
- Macoma nasuta* (Conrad)
 Grant and Gale, 1931, p. 365, pl. 20, fig. 11a, 11b.
 Hypotype; 178; Pleistocene; southwest of Goleta, Santa Barbara Co., Calif.
- Macoma panzana* Wiedey
 Wiedey, 1928, p. 150, pl. 19, fig. 1.
 Holotype; 38; Miocene, Temblor Formation; 2 miles west of Simmler, San Luis Obispo Co., Calif.
- Macoma secta* (Conrad)
 Grant and Gale, 1931, p. 374, pl. 20, fig. 6a, 6b.
 Hypotype; 177; Pliocene, Pico Formation; Holser Canyon, Los Angeles Co., Calif.

Macoma sespeensis Loel and Corey

Loel and Corey, 1932, p. 228.

Paratype; 789; Miocene, Vaqueros Formation; one mile up Little Sespe Creek from junction with Sespe Creek, Ventura Co., Calif.

Macrocallista stantoni Waring

Waring, 1917, p. 77.

Paratype; 623; Eocene, Martinez Formation; Martinez area, Simi Hills, Ventura Co., Calif.

Mactra (Mactra) orthomorpha Grant and Gale

Grant and Gale, 1931, p. 391, pl. 23, fig. 2a-2c, 6, 7.

Holotype; 193 (fig. 2a-2c); paratype; 194 (fig. 6), 195 (fig. 7), 746 (fig. 2b, pars, "fragment of another specimen"); Pliocene, Etchegoin Formation; oil well near Tipton, Tulare Co., Calif.

Mactra (Spisula) albaria Conrad

Grant and Gale, 1931, p. 395, pl. 23, fig. 3a, 3b.

Hypotype; 189; Miocene; 4575 feet deep in oil well northwest of Bakersfield, Kern Co., Calif.

Mactra (Spisula) catilliformis (Conrad)

Grant and Gale, 1931, p. 398, pl. 23, fig. 4, 10.

Hypotype; 196 (fig. 4), 197 (fig. 10); Pliocene, Etchegoin Formation; oil well 3229-3230 feet deep, northwest of Bakersfield, Kern Co., Calif.

Mya (Mya) arenaria Linnaeus variety *japonica* Jay

Grant and Gale, 1931, p. 412, pl. 21, fig. 13.

Hypotype; 188; Pliocene, Etchegoin Formation; 2877 feet deep in oil well, Kern Co., Calif.

Mya (Platyonodon) cancellata Conrad

Grant and Gale, 1931, p. 415, pl. 24, fig. 3a, 3b.

Hypotype; 202; Pleistocene; southwest of Goleta, Santa Barbara Co., Calif.

Nucula (Acula) sp. Grant and Gale

Grant and Gale, 1931, p. 115, text-fig. 4.

Hypotype; 374; Pliocene; 1 1/4 miles west of San Martinez Grande Canyon, Los Angeles Co. and Ventura Co., Calif., boundary line.

Nucula (Acula) gettysburgensis Reagan

Grant and Gale, 1931, p. 113, text-fig. 1.

Hypotype; 375; Oligocene; Twin Rivers Shales; one mile west of the mouth of West Twin River, Clallam Co., Wash.

Nucula (Acula) semirostrata Grant and Gale

Grant and Gale, 1931, p. 113, text-fig. 2a, 2b, 3a, 3b.

Holotype; 370 (fig. 2a, 2b); paratype; 371 (fig. 3a, 3b), 372, 373; Pliocene; 1 1/4 miles west of San Martinez Grande Canyon, Los Angeles Co. and Ventura Co., Calif., boundary line.

Nucula (Ennucula) birchi Keen

Keen, 1943, p. 41.

Paratype; 611, 933; Miocene, Round Mountain Silt; Caliente Quadrangle, Kern Co., Calif.

Nucula (Ennucula) microsperma Berry

Berry, 1947, p. 258 (4).

Paratype; 318, 319; Pleistocene, Lomita Formation; 2nd and Pacific Streets, San Pedro, Los Angeles Co., Calif.

Nuculana taphria (Dall)

Grant and Gale, 1931, p. 121, pl. 1, fig. 8, 9.

Hypotype; 77 (fig. 8), 78 (fig. 9); 77: Pliocene; Holser Canyon, Los Angeles Co., Calif.; 78: Pleistocene; near Goleta, Santa Barbara Co., Calif.

Ostrea angelica Rochebrune

Emerson and Hertlein, 1964, p. 353, fig. 3a, 3b.

Hypotype; 802; ?Pliocene; Isla Angel de la Guarda, Baja California, Mexico.

Ostrea ashleyi Hertlein

Hertlein, 1934, p. 1.

Paratype; 429; Miocene, Tremblor Formation; Kern Co., Calif.

Ostrea eldridgei (Arnold) *ynezana* Loel and Corey

Loel and Corey, 1932, p. 189.

Paratype; 779; Miocene, Vaqueros Formation; El Jaro Creek, Santa Ynez Mountains, Santa Barbara Co., Calif.

Ostrea haleyi Hertlein

(Hertlein, 1933, p. 277, pl. 18, fig. 5, 6.)

Plastoholotype; 425 (fig. 5), 426 (fig. 6); paratype 427; Eocene; Santa Cruz Island, Calif.

Ostrea howelli Wiedey

Wiedey, 1928, p. 135, pl. 15, fig. 1, 2.

Holotype; 26; Miocene; 5 1/2 miles northeast of Wheeler's Hot Springs, Ventura Co., Calif.

Ostrea tayloriana Gabb

(Gabb, 1869, p. 34, pl. 12, fig. 60, 60a.)

Plastoholotype; 190, 742; Eocene; San Marcos Pass, Santa Barbara Co., Calif.

Ostrea vaquerosensis Loel and Corey

Loel and Corey, 1932, p. 192.

Paratype; 780; Miocene, Vaqueros Formation; 2.6 miles north 21 degrees east of Abalone Point, Orange Co., Calif.

Ostrea venturana Loel and Corey

Loel and Corey, 1932, p. 193.

Paratype; 781; Miocene, Vaqueros Formation; ridge north of Coyote Creek fork of Ventura River, Ventura Co., Calif.

Ostrea vespertina Conrad

Grant and Gale, 1931, p. 152, pl. 12, fig. 1a, 1b.

Hypotype; 116; Pliocene, Pico Formation; Holser Canyon, Los Angeles Co., Calif.

Pandora punctata Conrad

Grant and Gale, 1931, p. 262, pl. 13, fig. 2a, 2b.

Hypotype; 121; Pleistocene, San Pedro Formation; Palos Verdes, Los Angeles Co., Calif.

Panope (Panope) generosa Gould

Grant and Gale, 1931, p. 424, pl. 21, fig. 12a, 12b.

Hypotype; 187; Pleistocene; southwest of Goleta, Santa Barbara Co., Calif.

Panope tenuis Wiedey

Wiedey, 1928, p. 154, pl. 20, fig. 4.

Holotype; 41; Miocene, Temblor Formation; 2 miles west of Simmler, San Luis Obispo Co., Calif.

Paphia restorationensis Frizzell

Frizzell, 1930, p. 120.

Holotype; 386; Pleistocene; near Port Blakely, Jefferson Co., Wash.

See: *Venerupis (Protobraca) restorationensis* (Frizzell)

Parallelodon (Nanonavis) vancouverensis (Meek)

Reinhart, 1937, p. 171.

See: *Arca vancouverensis* Meek

Pecten (Aequipecten) andersoni Arnold

Grant and Gale, 1931, p. 202, pl. 4, fig. 4, 5.

Hypotype; 93 (fig. 4), 94 (fig. 5); Miocene; Barker's Ranch, Kern Co., Calif.

Pecten (Aequipecten) deserti Conrad, type variety

Grant and Gale, 1931, p. 212, pl. 5, fig. 3.

Hypotype; 98; Pliocene, Carrizo Creek beds, Imperial Co., Calif.

Pecten (Aequipecten) deserti Conrad variety *invalidus* Hanna

Grant and Gale, 1931, p. 213, pl. 5, fig. 5a-c, 6a-c.

Hypotype; 100 (fig. 5a-c), 101 (fig. 6a-c); Pliocene, Pico Formation; between Holser and San Martinez Grande Canyons, Los Angeles Co., Calif.

Pecten (Aequipecten) discus Conrad

Grant and Gale, 1931, p. 200, pl. 4, fig. 7.

Hypotype; 96; Miocene, Santa Margarita Formation; 4655 foot depth in Ansolabehere No. 1 oil well, northwest of Bakersfield, Kern Co., Calif.

Pecten (Aequipecten) latiauratus Conrad variety *monotimeris* Conrad

Grant and Gale, 1931, p. 204, pl. 4, fig. 3, 6.

Hypotype; 92 (fig. 3), 95 (fig. 6); Pleistocene; ½ mile south of Southern Pacific station at Seadliff, Ventura Co., Calif.

Pecten (Aequipecten) percarus Hertlein

Hertlein, 1925a, p. 13.

Paratype; 630; Pliocene, Salada Formation; northwest of Elephant Mesa west of Arroyo, Scammons Lagoon Quadrangle, Baja California, Mexico.

Pecten (Aequipecten) purpuratus Lamarck variety *callidus* Hertlein

Grant and Gale, 1931, p. 211, pl. 5, fig. 4.

Hypotype; 99; Pliocene, Pico Formation; west of Fernando Pass, Los Angeles Co., Calif.

Pecten (Aequipecten) purpuratus Lamarck variety *subdolus* Hertlein

Grant and Gale, 1931, p. 211, pl. 5, fig. 1.

Hypotype; 97; Pliocene, Pico Formation; summit of ridge over Fernando Pass railroad tunnel, Los Angeles Co., Calif.

Pecten (Camptonectes) harfordus Davis

Davis, 1913, p. 456.

Paratype; 634, 938, 939; Jurassic, San Luis Formation; 6 miles north of Port Harford, San Luis Obispo Co., Calif.

Pecten (Chlamys) crici Wiedey

Wiedey, 1928, p. 137, pl. 16, fig. 1.

Holotype; 27; Miocene, Vaqueros Formation; South Mountain, Ventura Co., Calif.

Pecten (Chlamys) hodgei Hertlein

Hertlein, 1925b, p. 42.

Paratype; 632; Miocene, Santa Margarita Formation; Coalinga region, Fresno Co., Calif.

Pecten (Janira) bellus (Conrad) variety *coalingensis* Arnold

Grant and Gale, 1931, p. 227, pl. 2, fig. 2.

Hypotype; 86; Pliocene, Pico Formation; southeast of Pico Canyon, Los Angeles Co., Calif.

Pecten (Janira) bellus (Conrad) variety *hemphilli* Dall

Grant and Gale, 1931, p. 226, pl. 3, fig. 1a, 1b.

Hypotype; 88a (fig. 1a), 88b (fig. 1b); Pliocene, San Diego Formation; Pacific Beach, San Diego, San Diego Co., Calif.

Pecten (Janira) bellus (Conrad) variety *slevini* Dall and Ochsner

Grant and Gale, 1931, p. 227, pl. 2, fig. 3.

Hypotype; 87; Pliocene; north of Santa Clara Valley and east of San Martinez Chiquito Canyon, Los Angeles Co., Calif.

Pecten (Janira) stearnsii Dall variety *bakeri* Hanna and Hertlein

Grant and Gale, 1931, p. 224, pl. 4, fig. 1a, 1b.

Hypotype; 91; Pliocene, Santa Rosalia, Territorio de Sur de Baja California, Mexico.

Pecten (Janira) stearnsii Dall variety *diegensis* Dall

Grant and Gale, 1931, p. 223, pl. 3, fig. 4.

Hypotype; 89; Pliocene, Pico Formation?; Smith Canyon, Ventura Co., Calif.

Pecten (Janira) stearnsii Dall variety *stearnsii*, s. s.

Grant and Gale, 1931, p. 223, pl. 3, fig. 2b (not 2a).

Hypotype; 90; Pliocene, Pico Formation; Holser Canyon, Los Angeles Co., Calif.

Pecten (Lyropecten) estrellanus (Conrad)

Grant and Gale, 1931, p. 185, pl. 8, fig. 4.

Hypotype; 106; Pliocene, Pico Formation; Elsmere Canyon, Los Angeles Co., Calif.

Pecten (Lyropecten) estrellanus (Conrad) variety *cerrosensis* Gabb

Grant and Gale, 1931, p. 187, pl. 8, fig. 1a, 1b, 2a, 2b, pl. 9, fig. 2.

Hypotype; 107 (pl. 8, fig. 1a, 1b), 108 (pl. 8, fig. 2a, 2b), 109 (pl. 9, fig. 2); Pliocene, Los Angeles Co., Calif.; 107: Pico Formation, southeast of Pico Canyon; 108: Pico Formation, Holser Canyon; 109: ridge north of Junction of Tomsley and Wiley Canyons.

Pecten (Pallium) swiftii Bernardi

Grant and Gale, 1931, p. 171, pl. 10, fig. 2, 5.

Hypotype; 110 (fig. 2), 113 (fig. 5); Pliocene, Pico Formation, Holser Canyon; Los Angeles Co., Calif.

Pecten (Pallium) swiftii Bernardi variety *etchegeoroi* Anderson

Grant and Gale, 1931, p. 173, pl. 10, fig. 3a, 3b.

Hypotype; 111; Pliocene, Purisima Formation; Purisima Creek, San Mateo Co., Calif.

Pecten (Patinopecten) caurinus Gould

Grant and Gale, 1931, p. 194, pl. 6, fig. 4.

Hypotype; 105; Pliocene, Purisima Formation; Purisima Creek, San Mateo Co., Calif.

Pecten (Patinopecten) healeyi Arnold

Grant and Gale, 1931, p. 196, pl. 6, fig. 2a, 2b.

Hypotype; 103a (fig. 2a), 103b (fig. 2b); Pliocene, San Diego Formation; Pacific Beach, San Diego, San Diego Co., Calif.

Pecten (Patinopecten) healeyi Arnold variety *lobri* Hertlein

Grant and Gale, 1931, p. 197, pl. 6, fig. 1a, 1b.

Hypotype; 102; Pliocene, Pico Formation; Elsmere Canyon, Los Angeles Co., Calif.

Pecten (Patinopecten) marquerensis (Durham)

Emerson and Hertlein, 1964, fig. 4f.

Hypotype; 810; Pliocene; Isla San José, Baja California, Mexico.

Pecten (Patinopecten) purisimaeensis Arnold

Grant and Gale, 1931, p. 194, pl. 6, fig. 3.

Hypotype; 104; Pliocene, Purisima Formation; mouth of San Gregorio Creek, San Mateo Co., Calif.

Pecten (Pecten) beringianus Middendorff

Grant and Gale, 1931, p. 165, pl. 11, fig. 2.

Hypotype; 114; Pliocene, Pico Formation; Sulphur Canyon, Ventura Co., Calif.

Pecten (Pecten) islandicus Muller variety *jordani* Arnold

Grant and Gale, 1931, p. 164, pl. 11, fig. 4.

Hypotype; 115; Pliocene; Packard's Hill, Santa Barbara, Santa Barbara Co., Calif.

Pecten (Plagipectenium) cristobalensis Hertlein

Hertlein, 1925a, p. 19.

Paratype; 631; Pliocene, Salada Formation; 3 miles southeast of Turtle Bay, San Cristobal Bay Quadrangle, Baja California, Mexico.

Pecten (Plagipectenium) hakei Hertlein

Hertlein, 1925a, p. 18.

Paratype; 638; Pliocene; Ballenas Quadrangle, Baja California, Mexico.

Pecten (Pseudantusium) vancouverensis fernandoensis Hertlein

Hertlein, 1925b, p. 43.

Paratype; 635; Pliocene, Fernando Formation; on Ventura River, 1.5 miles north of Ventura, Ventura Co., Calif.

Pecten vanwinkleae Clark

Clark, 1925, p. 82.

Paratype; 636; Oligocene, Lincoln Formation; along Porter Creek about $\frac{3}{4}$ miles above Porter, Grays Harbor Co., Wash.

Pedion panzana Loel and Corey

Loel and Corey, 1932, p. 187.

Paratype; 778; Miocene, Vaqueros Formation; branch of Carrizo Creek about 100 yards west of Carrizo ranch house, La Panza Mountains, San Luis Obispo Co., Calif.

Petricola carditoides (Conrad)

Grant and Gale, 1931, p. 355, pl. 13, fig. 14a, 14b.

Hypotype; 128; Pleistocene; southwest of Goleta, Santa Barbara Co., Calif.

Pholadidea (Pholadidea) penita (Conrad)

Grant and Gale, 1931, p. 434, pl. 24, fig. 1a, 1b.

Hypotype; 200; Pleistocene; southwest of Goleta, Santa Barbara Co., Calif.

Pholadomya kernensis Wiedey

(Wiedey, 1928, p. 141, pl. 17, fig. 1, 2.)

Plastoholotype; 191, 743-745; Miocene, Temblor Formation; north of Poso Creek, Kern Co., Calif.

Pholas (Zirfaea) gabbi (Tryon)

Grant and Gale, 1931, p. 432, pl. 24, fig. 2.

Hypotype; 201; Pliocene, Etchegoin Formation; 3900-18 feet deep in oil well near McFarland, Kern Co., Calif.

Pinna latraria Hanna

Hanna, 1926, p. 476.

Paratype; 1; Pliocene, Imperial Formation; Coyote Mountain, Imperial Co., Calif.

Pododesmus macroschisma (Deshayes)

Grant and Gale, 1931, p. 241, pl. 12, fig. 3.

Hypotype; 120; Pliocene, Pico Formation; Holser Canyon, Los Angeles Co., Calif.

Pteria jordani Wiedey

(Wiedey, 1928, p. 134, pl. 14, fig. 4.)

Holotype; 25; Miocene, Temblor Formation; 2 miles south of Calabasas, Los Angeles Co., Calif.

Sanguinolaria (Nuttallia) orcutti Dall

Dall, 1921, p. 17.

Syntype; 940-950; Pleistocene; San Quintin Bay, Baja California, Mexico.

Saxidomus nuttalli Conrad variety *giganteus* (Deshayes)

Grant and Gale, 1931, p. 342, pl. 18, fig. 4, 10.

Hypotype; 156 (fig. 4), 158 (fig. 10); 156: Pleistocene, 1 to $1\frac{1}{4}$ miles west of Goleta Point, Santa Barbara Co., Calif.; 158: Pliocene, 100 yards west of the bathhouse, Santa Barbara, Santa Barbara Co., Calif.

Schizothaerus nuttallii (Conrad)

Grant and Gale, 1931, p. 404, pl. 22, fig. 6a, 6b, pl. 23, fig. 8a, 8b, 9.

Hypotype; 198 (pl. 23, fig. 8a, 8b), 199 (pl. 23, fig. 9), 330 (pl. 22, fig. 6a, 6b); 198, 199: Pleistocene, southwest of Goleta, Santa Barbara Co., Calif.; 330: Pliocene, Purisima Formation, Pt. Año Nuevo, San Mateo Co., Calif.

Semele decis (Conrad)

Grant and Gale, 1931, p. 376, pl. 14, fig. 13a, 13b.

Hypotype; 137; Pleistocene, San Pedro Formation; San Pedro, Los Angeles Co., Calif.

Semele verrucosa Mörch

Emerson and Hertlein, 1964, p. 359, fig. 3i, 3j.

Hypotype; 803; Pleistocene; Isla Coronados, Baja California, Mexico.

Siliqua cf. patula (Dixon)

Grant and Gale, 1931, p. 924, pl. 21, fig. 9.

Hypotype; 186; Pliocene, Etchegoin Formation; oil well near McFarland, Kern Co., Calif.

Siliqua lucida (Conrad)

Grant and Gale, 1931, p. 389, pl. 21, fig. 6.

Hypotype; 184; Pliocene, Etchegoin Formation; 3000-18 feet deep in well near McFarland, Kern Co., Calif.

Solen sicarius Gould

Grant and Gale, 1931, p. 385, pl. 21, fig. 4.

Hypotype; 183; Pleistocene; southwest of Goleta, Santa Barbara Co., Calif.

Sphaerium (Amesoda) rogersi Hannibal

Hannibal, 1913, p. 131.

Paratype; 633; Eocene, Tejon Formation; $\frac{1}{4}$ mile above Carnegie Pottery in Western Pacific Railway cut, Corral Hollow, Tesla, Alameda Co., Calif.

Spisula abbotti Wiedey

Wiedey, 1928, p. 151, pl. 19, fig. 4, 5.

Holotype; 39; Miocene, Temblor Formation; 2 miles northwest of powerhouse at mouth of Kern Canyon, Kern Co., Calif.

Spisula granti Wiedey

Wiedey, 1928, p. 152, pl. 20, fig. 2, 3.

Holotype; 40; Miocene, Vaqueros Formation; Vaqueros Creek, Monterey Co., Calif.

Spondylus inezana Wiedey

Wiedey, 1928, p. 139, pl. 16, fig. 2, 3.

Holotype; 29; Miocene, Vaqueros Formation; Head of Wiley Canyon, Ventura Co., Calif.

Spondylus perrini Wiedey

Wiedey, 1928, p. 138, pl. 17, fig. 6, 7.

Holotype; 28; Miocene, Vaqueros Formation; Head of Wiley Canyon, Ventura Co., Calif.

Tagelus californianus (Conrad)

Grant and Gale, 1931, p. 384, pl. 21, fig. 2a, 2b, 3.

Hypotype; 181 (fig. 2a, 2b), 182 (fig. 3); Pleistocene; San Pedro, Los Angeles Co., Calif.

Tellina bodegaensis Hinds

Grant and Gale, 1931, p. 362, pl. 20, fig. 13.

Hypotype; 180; Pleistocene; west of Newport, Los Angeles Co., Calif.

Tellina idae Dall

Grant and Gale, 1931, p. 358, pl. 20, fig. 12.

Hypotype; 179; Pliocene, Pico Formation; above Newhall railroad tunnel, Los Angeles Co., Calif.

Tellina oldroydi Wiedey

Wiedey, 1928, p. 148, pl. 19, fig. 3.

Holotype; 36; Miocene, Temblor Formation; 2 miles south of Calabasas, Los Angeles Co., Calif.

Thracia (Thracia) trapezoides Conrad

Grant and Gale, 1931, p. 257, pl. 13, fig. 8.

Hypotype; 123; Miocene; 2 or 3 miles south of Mayfield, Santa Clara Co., Calif.

Thyasira bisecta (Conrad) variety *nipponica* Yabe and Nomura

Grant and Gale, 1931, pl. 13, fig. 15.

Hypotype; 129; Pliocene or Pleistocene, Kawabata Series; Japan.

Tivela gasterensis Clark

Clark, 1925, p. 93.

Paratype; 627; Oligocene; county quarry, Scroggins Canyon, Gaston, Washington Co., Oregon.

Tivela stultorum (Mawe)

Grant and Gale, 1931, p. 340, pl. 19, fig. 3a, 3b.

Hypotype; 170; Pleistocene; west of Newport, Los Angeles Co., Calif.

Venerupis (Protobrachia) restorationensis (Frizzell)

Frizzell, 1931, p. 321, pl. 22, fig. 2-4.

Holotype; 386; Pleistocene; near Port Blakely, Jefferson Co., Wash.

See: *Paphia restorationensis* Frizzell

Venerupis (Protobrachia) staminea (Conrad)

Grant and Gale, 1931, p. 329, pl. 18, fig. 1a, 1b, 2a, 2b.

Hypotype; 153 (fig. 1a, 1b), 154 (fig. 2a, 2b); Pleistocene; beach southwest of Goleta, Santa Barbara Co., Calif.

Venerupis (Protobrachia) staminea (Conrad) variety *ruderata* (Deshayes)

Grant and Gale, 1931, p. 331, pl. 18, fig. 3a, 3b.

Hypotype; 155; Pleistocene; 1 to 1 1/4 miles west of Goleta Point, Santa Barbara Co., Calif.

Venus (Chione) elsmereensis (English)

Grant and Gale, 1931, p. 319, pl. 16, fig. 6a, 6b, 7.

Hypotype; 151 (fig. 6a, 6b), 152 (fig. 7); Pliocene, Pico Formation; Elsmere Canyon, Los Angeles Co., Calif.

Venus (Chione) securis Shumard variety *fernandoensis* (English)

Grant and Gale, 1931, p. 321, pl. 17, fig. 4a, 4b, 5, 6.

Hypotype; 159 (fig. 4a, 4b), 160 (fig. 5), 161 (fig. 6); Pliocene, Pico Formation; Elsmere Canyon, Los Angeles Co., Calif.

Venus (Chione) succincta Valenciennes

Grant and Gale, 1931, p. 321, pl. 16, fig. 1a, 1b, 2a, 2b, 3, 4.

Hypotype; 147 (fig. 1a, 1b), 148 (fig. 2a, 2b), 149 (fig. 3), 150 (fig. 4); Pleistocene; 5 miles ENE of Newport Beach, Orange Co., Calif.

Gastropoda

Acanthina spirata (Blainville)

Grant and Gale, 1931, p. 720, pl. 32, fig. 6, 7.

Hypotype; 293 (fig. 6), 294 (fig. 7); Pleistocene; 293: San Pedro, Los Angeles Co., Calif.; 294: Sexton Canyon, Ventura Co., Calif.

Acmaea instabilis (Gould)

Grant and Gale, 1931, p. 813, pl. 32, fig. 32.

Hypotype; 313; Pleistocene; southwest of Goleta, Santa Barbara Co., Calif.

Acmaea lepsima Berry

Berry, 1940, p. 9.

Paratype; 353, 354, 355; Pleistocene; Hilltop Quarry, San Pedro, Los Angeles Co., Calif.

Acteon (Rictaxis) painei Dall variety *grandior* Grant and Gale

Grant and Gale, 1931, p. 444, pl. 24, fig. 12.

Holotype; 208; Pliocene, Pico Formation; Holser Canyon, Los Angeles Co., Calif.

Admete modesta (Carpenter)

Grant and Gale, 1931, p. 622, pl. 27, fig. 5.

Hypotype; 267; Pliocene, Pico Formation; Sulphur Canyon, Ventura Co., Calif.

Amphissa columbiana Dall

Grant and Gale, 1931, p. 701, pl. 26, fig. 39.

Hypotype; 250; Pleistocene; southwest of Goleta, Santa Barbara Co., Calif.

Amphissa versicolor Dall

Grant and Gale, 1931, p. 702, pl. 26, fig. 53.

Hypotype; 259; Pleistocene; California State Highway 1500 feet north of Rincon Creek, Santa Barbara Co., Calif.

Architectonica compressa Wiedey

Wiedey, 1928, p. 109, pl. 9, fig. 1, 2.

Holotype; 12; Miocene, Temblor Formation; 2 miles southeast of El Modena, Orange Co., Calif.

Architectonica nobilis Bolten variety *discus* Grant and Gale

Grant and Gale, 1931, p. 786, pl. 32, fig. 27.

Holotype; 308; Pliocene?; one mile southwest of Yuha drill hole, south of Dixieland, Imperial Co., Calif.

Astrea (Pachypoma) inaequalis (Martyn)

Grant and Gale, 1931, p. 820, pl. 31, fig. 4a, 4b.

Hypotype; 284; Pliocene; Bath-House Beach, Santa Barbara, Santa Barbara Co., Calif.

Astrea (Pomaularax) gradata Grant and Gale

Grant and Gale, 1931, p. 818, pl. 31, fig. 1a, 1b, 3a, 3b, 5, 8, 9.

Holotype; 286 (fig. 8); paratype; 282 (fig. 1a, 1b), 283 (fig. 3a, 3b), 285 (fig. 5), 287 (fig. 9); Pliocene, Pico Formation; 282, 285, 287; Holser Canyon; 283: west of Fernando Pass; 286: between Pico Canyon and Fernando Pass; all Los Angeles Co., Calif.

Balcis (Balcis) clavella Berry

Berry, 1954, p. 5.

Paratype; 413; Pleistocene; Long Wharf Canyon, Santa Monica, Los Angeles Co., Calif.

Balcis (Balcis) tersa Berry

Berry, 1954, p. 7.

Paratype; 395, 396, 397; Pleistocene; "Hilltop Quarry", San Pedro, Los Angeles Co., Calif.

Balcis conchita Keen

Keen, 1943, p. 43.

Paratype; 617, 936; Miocene, Round Mountain Silt; Caliente Quadrangle, Kern Co., Calif.

Balcis (Vitreolina) ebrikanus Berry

Berry, 1954, p. 11.

Paratype; 410, 411, 412; Pleistocene; "Hilltop Quarry", San Pedro, Los Angeles Co., Calif.

Balcis (Vitreolina) incallida Berry

Berry, 1954, p. 10.

Paratype; 404-409; Pliocene; "Hilltop Quarry", San Pedro, Los Angeles Co., Calif.

Balcis (Vitreolina) obstipa Berry

Berry, 1954, p. 8.

Paratypes; 398, 399, 400, 401, 402, 403; Pleistocene; "Hilltop Quarry", San Pedro, Los Angeles Co., Calif.

Bathytoma clarkiana Rivers

Rivers, 1913, p. 29, 2 figures on unnumbered plate.

Syntypes; 10 (figured on left side of plate), 11 (figured on right side of plate); age and locality uncertain.

Bitium (Lirobitium) asperum (Gabb) variety *dilatatum* Grant and Gale

Grant and Gale, 1931, p. 760, pl. 24, fig. 14.

Holotype; 210; Pliocene, Pico Formation; Holser Canyon, Los Angeles Co., Calif.

Bitium (Semibitium) rugatum Carpenter

Grant and Gale, 1931, p. 762, pl. 24, fig. 8.

Hypotype; 205; Pleistocene; southwest of Goleta, Santa Barbara Co., Calif.

Calliostoma canaliculatum (Martyn)

Grant and Gale, 1931, p. 833, pl. 32, fig. 23.

Hypotype; 306; Pleistocene; San Pedro, Los Angeles Co., Calif.

- Callistoma gemmulum* Carpenter
 Grant and Gale, 1931, p. 835, pl. 32, fig. 21.
 Hypotype; 305; Pleistocene; one half mile south of Seacliff Station, Ventura Co., Calif.
- Calyptraea mammillaria* Broderip
 Grant and Gale, 1931, p. 794, pl. 32, fig. 24a, 24b.
 Hypotype; 307; Pleistocene; southwest of Goleta, Santa Barbara Co., Calif.
- Cancellaria clavatula* Sowerby
 Grant and Gale, 1931, p. 615, pl. 27, fig. 2.
 Hypotype; 264; Pliocene, Pico Formation; east of Fernando Pass, Los Angeles Co., Calif.
- Cancellaria hemphilli* Dall
 Grant and Gale, 1931, p. 621, pl. 27, fig. 3, 15a, 15b.
 Hypotype; 265 (fig. 3), 272 (fig. 15a, 15b); Pliocene; 265: Pico Formation, Holser Canyon, Los Angeles Co., Calif.; 272: between Pico Canyon and Fernando Pass, Los Angeles Co., Calif.
- Cancellaria obesa* Sowerby variety *planospira* Grant and Gale
 Grant and Gale, 1931, p. 613, pl. 27, fig. 4.
 Holotype; 266; Pliocene, Pico Formation; east of Fernando Pass, Los Angeles Co., Calif.
- Cancellaria tritonidea* Gabb variety *fernandoensis* Arnold
 Grant and Gale, 1931, p. 618, pl. 27, fig. 1.
 Hypotype; 263; Pliocene, Pico Formation; east of Fernando Pass, Los Angeles Co., Calif.
- Cantharus fortis* (Carpenter)
 Grant and Gale, 1931, p. 647, pl. 28, fig. 2.
 Hypotype; 273; Pleistocene; Santa Paula Creek, Ventura Co., Calif.
- Cantharus humerosus* (Gabb)
 Grant and Gale, 1931, p. 647, pl. 28, fig. 3.
 Hypotype; 274; Pliocene; between Pico Canyon and Fernando Pass, Los Angeles Co., Calif.
- Cerithidea californica* (Haldeman)
 Grant and Gale, 1931, p. 763, pl. 24, fig. 6, 7.
 Hypotype; 203 (fig. 6), 204 (fig. 7); Pleistocene; San Pedro, Los Angeles Co., Calif.
- "*Cerithium*" *simplicius* Grant and Gale
 Grant and Gale, 1931, p. 757, pl. 24, fig. 11.
 Holotype; 207; Pliocene, Pico Formation; Holser Canyon, Los Angeles Co., Calif.
- Chicoreus (Murithais) wilkesanus* (Anderson)
 Grant and Gale, 1931, p. 730, pl. 32, fig. 12.
 Hypotype; 298; Miocene; oil well northwest of Bakersfield, Kern Co., Calif.
- Chrysallida rotundomontana* Keen
 Keen, 1943, p. 43.
 Paratype; 610; Miocene, Round Mountain Silt; Caliente Quadrangle, Kern Co., Calif.
- Clathurella coradiana* Gabb
 Grant and Gale, 1931, p. 606, pl. 26, fig. 11.
 Hypotype; 236; Pliocene, Pico Formation; Sulphur Canyon, Ventura Co., Calif.
- Clathurella (Glyphostoma) tridesmia* Berry
 Berry, 1941, p. 8.
 Paratype; 384, 385, 393; Pleistocene; "Hilltop Quarry"; San Pedro, Los Angeles Co., Calif.
- Clavus (Clathrodrillia) coalingensis* (Arnold)
 Grant and Gale, 1931, p. 580, pl. 26, fig. 14a, 14b, 15.
 Hypotype; 232 (fig. 14a, 14b), 233 (fig. 15); Pliocene, Pico Formation; Elsmere Canyon, Los Angeles Co., Calif.
- Clavus (Crassispira)* sp.
 Grant and Gale, 1931, p. 582, pl. 26, fig. 10.
 Hypotype; 231; Pliocene, Pico Formation; Holser Canyon, Los Angeles Co., Calif.
- Clavus (Cymatosyrinx) pallidus* (Sowerby)
 Grant and Gale, 1931, p. 576, pl. 26, fig. 16a, 16b, 17.
 Hypotype; 234 (fig. 16a, 16b), 235 (fig. 17); Pliocene, Pico Formation; Holser Canyon, Los Angeles Co., Calif.
- Conus californicus* Hinds
 Grant and Gale, 1931, p. 472, pl. 24, fig. 21.
 Hypotype; 216; Pleistocene; Santa Paula Quadrangle, Ventura Co., Calif.
- Conus californicus fossilis* Oldroyd
 Oldroyd, T. S., 1921, p. 116.
 Paratype; 361, 362; Pleistocene; San Pedro, Los Angeles Co., Calif.
- Conus juanensis* Wiedey
 Wiedey, 1928, p. 123, pl. 9, fig. 3.
 Holotype; 16; Miocene, Temblor Formation; 2 miles west of Simmler, San Luis Obispo Co., Calif.
- Conus Oweniana* (Anderson) *ynezanus* Loel and Corey
 Loel and Corey, 1932, p. 237.
 Paratype; 794; Miocene, Vaqueros Formation; 2.5 miles southwest of Buellton, Santa Barbara Co., Calif.

- Crepidula onyx* Sowerby
 Grant and Gale, 1931, p. 790, pl. 32, fig. 34.
 Hypotype; 314; Pliocene, Pico Formation; Canada de Aliso, Ventura Co., Calif.
- Cyllichna* ? *loismartinae* Keen
 Keen, 1943, p. 44.
 Paratype; 608; Miocene, Round Mountain Silt; Caliente Quadrangle, Kern Co., Calif.
- Cyllichna temblorensis* Keen
 Keen, 1943, p. 44.
 Paratype; 609, 930, 931, 932; Miocene, Round Mountain Silt; Caliente Quadrangle, Kern Co., Calif.
- Cypraea spadicea* Swainson
 Grant and Gale, 1931, p. 752, pl. 27, fig. 13.
 Hypotype; 270; Pliocene, Pico Formation; Holser Canyon, Los Angeles Co., Calif.
- Drillia merriami* Arnold
 (Arnold, 1903, p. 207, pl. 8, fig. 7.)
 Plastoholotype; 637; Pleistocene, San Pedro Formation; Deadman Island, San Pedro, Los Angeles Co., Calif.
- Epitonium* (*Opalia*) *varicostatum* (Stearns)
 Grant and Gale, 1931, p. 853, pl. 24, fig. 20.
 Hypotype; 215; Pliocene, San Diego Formation; Pacific Beach, San Diego Co., Calif.
- Ferminoscalra whitei* Keen
 Keen, 1943, p. 46.
 Paratype; 607, 929; Miocene, Round Mountain Silt; Caliente Quadrangle, Kern Co., Calif.
- Ficus* (*Trophoscyon*) *ocoyana* (Conrad)
 Grant and Gale, 1931, p. 743, pl. 30, fig. 11.
 Hypotype; 281; Miocene; southwest of Mayfield, Santa Clara Co., Calif.
- Ficus* (*Trophoscyon*) *ocoyana* (Conrad) variety *contignata* Grant and Gale
 Grant and Gale, 1931, p. 749, pl. 29, fig. 1a, 1b, pl. 30, fig. 2, 4, 9a, 9b.
 Holotype; 4 (pl. 29, fig. 1a, 1b, pl. 30, fig. 2); paratype; 278 (pl. 30, fig. 4), 279 (pl. 30, fig. 9a, 9b); Pliocene, Pico Formation; 4, 279; Elsmere Canyon; 278: south of Humphrey's Station; both Los Angeles Co., Calif.
- Ficus* (*Trophoscyon*) *ocoyana* (Conrad) variety *ruginodosa* Grant and Gale
 Grant and Gale, 1931, p. 746, pl. 29, fig. 2a, 2b, pl. 30, fig. 10a, 10b.
 Holotype; 3 (pl. 29, fig. 2a, 2b); paratype (pl. 30, fig. 10a, 10b); Pliocene, Pico Formation; Elsmere Canyon, Los Angeles Co., Calif.
- Forrieria magister* (Nomland)
 Grant and Gale, 1931, p. 727, pl. 27, fig. 14a, 14b.
 Hypotype; 271; Pliocene, Pico Formation; Elsmere Canyon, Los Angeles Co., Calif.
- Fusinus barbarensis* (Trask)
 Grant and Gale, 1931, p. 639, pl. 27, fig. 11.
 Hypotype; 268; Pliocene; between Pico Canyon and Fernando Pass, Los Angeles Co., Calif.
- Galeodea apta* Tegland
 Tegland, 1931, p. 415.
 Paratype; 366, 367; Oligocene, Twin Rivers Shales; 1½ miles west of mouth of West Twin River, Clallam Co., Wash.
- Galeodea rex* Tegland
 Tegland, 1931, p. 413.
 Paratype; 363-365; Oligocene, Blakeley Formation; Bainbridge Island, Jefferson Co., Wash.
- Gyreneum* (*Bechtelia*) *strongi* Jordan
 Emerson and Hertlein, 1964, p. 360, fig. 5g.
 Hypotype; 818; Pleistocene; Isla Monserrate, Baja California, Mexico.
- Gyreneum lewisi* Carson
 Carson, 1926, p. 53.
 Paratype; 628; Pliocene, Fernando Formation; Fugler's Point, Santa Barbara Co., Calif.
- Hastula gnomon* Keen
 Keen, 1943, p. 47.
 Paratype; 606; Miocene, Round Mountain Silt; Caliente Quadrangle, Kern Co., Calif.
- Hyalina* (*Cystiscus*) *jewettii* (Carpenter)
 Grant and Gale, 1931, p. 630, pl. 24, fig. 17.
 Hypotype; 213; Pleistocene; San Pedro, Los Angeles Co., Calif.
- Kelletia* (*Kelletia*) *kelleteri* (Forbes)
 Grant and Gale, 1931, p. 642, pl. 28, fig. 7.
 Hypotype; 276; Pliocene, Pico Formation; Elsmere Canyon, Los Angeles Co., Calif.
- Lacuna divaricata* (Fabricius) variety *solidula* Lovén
 Grant and Gale, 1931, p. 782, pl. 32, fig. 19.
 Hypotype; 303; Pleistocene; southwest of Goleta, Santa Barbara Co., Calif.

- Littorina scutulata* Gould
 Grant and Gale, 1931, p. 782, pl. 32, fig. 16-18.
 Hypotype; 300 (fig. 16), 301 (fig. 17), 302 (fig. 18); Pleistocene; San Pedro, Los Angeles Co., Calif.
- Margarites (Lirularia) aresta* Berry
 Berry, 1941, p. 13.
 Paratype; 73; Pleistocene; "Hilltop Quarry", San Pedro, Los Angeles Co., Calif.
- Melampus olivaceous* Carpenter
 Grant and Gale, 1931, p. 461, pl. 24, fig. 16.
 Hypotype; 212; Pleistocene; San Pedro, Los Angeles Co., Calif.
- Miopleiona weaveri* Tegland
 (Tegland, 1933, p. 127, pl. 11, fig. 1, 2.)
 Plastoholotype; 360; Oligocene, Blakeley Formation; Bainbridge Island, Jefferson Co., Wash.
- Mistostigma punctulum* Berry
 Berry, 1947, p. 264 (10).
 Paratype; 322-324; Pliocene; Bath-house Cliff, Santa Barbara, Santa Barbara Co., Calif.
- Mitra catalinae* (Dall)
 Grant and Gale, 1931, p. 636, pl. 28, fig. 4.
 Hypotype; 275; Pliocene, Pico Formation; Holser Canyon, Los Angeles Co., Calif.
- Mitrella carinata* (Hinds)
 Grant and Gale, 1931, p. 692, pl. 26, fig. 35.
 Hypotype; 248; Pleistocene; ½ mile south of Sealiff Railway Station, Ventura Co., Calif.
- Mitrella carinata* (Hinds) variety *gausapata* (Gould)
 Grant and Gale, 1931, p. 693, pl. 26, fig. 44.
 Hypotype; 253; Pleistocene; south of Sealiff Railway Station, Ventura Co., Calif.
- Mitrella grandior* Grant and Gale
 Grant and Gale, 1931, p. 696, pl. 26, fig. 46.
 Holotype; 6; Pliocene, Pico Formation; Canada de Aliso, Ventura Co., Calif.
- Mitrella (Mitrella) anchuela* Keen
 Keen, 1943, p. 48.
 Paratype; 612; Miocene, Round Mountain Silt; Caliente Quadrangle, Kern Co., Calif.
- Mitrella tuberosa* (Carpenter)
 Grant and Gale, 1931, p. 697, pl. 26, fig. 45.
 Hypotype; 254; Pliocene, Pico Formation; Sulphur Canyon, Ventura Co., Calif.
- Mitromorpha galeana* Berry
 Berry, 1941, p. 12.
 Paratype; 54, 394; Pleistocene; "Hilltop Quarry", San Pedro, Los Angeles Co., Calif.
- Moniliopsis chacei* Berry
 Berry, 1941, p. 6.
 Paratype; 347, 348; Pleistocene; "Hilltop Quarry", San Pedro, Los Angeles Co., Calif.
- Moniliopsis graciosa* (Arnold) variety *mercedensis* (Martin)
 Grant and Gale, 1931, p. 569, pl. 26, fig. 30.
 Hypotype; 245; Pleistocene, Las Posas Formation; north of Arroyo Santa Rosa, Ventura Co., Calif.
- Moniliopsis incisa* (Carpenter) variety *incisa* s.s.
 Grant and Gale, 1931, p. 566, pl. 26, fig. 21.
 Hypotype; 238; Pleistocene, La Posas Formation; north of Arroyo Santa Rosa, Ventura Co., Calif.
- Moniliopsis incisa* (Carpenter) variety *quinquecincta* Grant and Gale
 Grant and Gale, 1931, p. 568, pl. 26, fig. 33.
 Hypotype; 247; Pliocene, Pico Formation; Elsmere Canyon, Los Angeles Co., Calif.
- Morula (Morunella) lugubris* (C. B. Adams)
 Emerson and Hertlein, 1964, p. 361, fig. 3k.
 Hypotype; 804; Pleistocene; Isla Coronados, Baja California, Mexico.
- Nassarius hildegardeae* Kanakoff
 Kanakoff, 1956, p. 113.
 Paratype; 341; Pliocene, Pico Formation; ½ mile south of Humphreys Railway Station, Los Angeles Co., Calif.
- Nassarius (Schizopyga) californianus* (Conrad)
 Grant and Gale, 1931, p. 672, pl. 26, fig. 49.
 Hypotype; 255; Pliocene, Pico Formation; Elsmere Canyon, Los Angeles Co., Calif.
 See: *Nassarius (Tritia) californianus* (Conrad)
- Nassarius (Schizopyga) fossatus* (Gould)
 Grant and Gale, 1931, p. 675, pl. 26, fig. 55, 56.
 Hypotype; 261 (fig. 55), 262 (fig. 56); Pleistocene; Southwest of Goleta, Santa Barbara Co., Calif.
 See: *Nassarius (Tritia) fossatus* (Gould)
- Nassarius (Schizopyga)mendicus* (Gould)
 Grant and Gale, 1931, p. 674, pl. 26, fig. 54.
 Hypotype; 260; Pleistocene; southwest of Goleta, Santa Barbara Co., Calif.

- See: *Nassarius (Tritia) mendicus* (Gould)
- Nassarius (Schizopyga) mendicus* (Gould) variety *cooperi* (Forbes)
Grant and Gale, 1931, p. 674, pl. 26, fig. 40, 50.
Hypotype; 251 (fig. 40), 256 (fig. 50); Pleistocene; southwest of Goleta, Santa Barbara Co., Calif.
- See: *Nassarius (Tritia) mendicus* (Gould) variety *cooperi* (Forbes)
- Nassarius (Schizopyga) perpinguis* (Hinds)
Grant and Gale, 1931, p. 673, pl. 26, fig. 51, 52.
Hypotype; 257 (fig. 51), 258 (fig. 52); 257; Pleistocene; southwest of Goleta, Santa Barbara Co., Calif.; 258; Pliocene, Pico Formation; Holser Canyon, Los Angeles Co., Calif.
- See: *Nassarius (Tritia) perpinguis* (Hinds)
- Nassarius (Tritia) californianus* (Conrad)
Grant and Gale, 1931, p. 941, pl. 26, fig. 49.
See: *Nassarius (Schizopyga) californianus* (Conrad)
- Nassarius (Tritia) fossatus* (Gould)
Grant and Gale, 1931, p. 675, pl. 26, fig. 55, 56.
See: *Nassarius (Schizopyga) fossatus* (Gould)
- Nassarius (Tritia) mendicus* (Gould)
Grant and Gale, 1931, p. 674, pl. 26, fig. 54.
See: *Nassarius (Schizopyga) mendicus* (Gould)
- Nassarius (Tritia) mendicus* (Gould) variety *cooperi* (Forbes)
Grant and Gale, 1931, p. 941, pl. 26, fig. 40, 50.
See: *Nassarius (Schizopyga) mendicus* (Gould) variety *cooperi* (Forbes)
- Nassarius (Tritia) perpinguis* (Hinds)
Grant and Gale, 1931, p. 673, pl. 26, fig. 51, 52.
See: *Nassarius (Schizopyga) perpinguis* (Hinds)
- Nassarius stocki* Kanakoff
Kanakoff, 1956, p. 110.
Paratype; 340; Pliocene, Pico Formation; 1/2 mile south of Humphreys Railway Station, Los Angeles Co., Calif.
- Nassarius (Uzita) arnoldi* (Anderson) variety *whitneyi* (Trask)
Grant and Gale, 1931, p. 679, pl. 26, fig. 48a, 48b.
Hypotype; 7; Pliocene, Etchegoin Formation; near Tipton, Tulare Co., Calif.
- Neptunea (Colus) jordani* (Dall)
Grant and Gale, 1931, p. 663, pl. 28, fig. 11.
Hypotype; 277; Pleistocene (Pliocene?), Elk River Formation; north of mouth of Elk River, Curry Co., Oregon.
- Neptunea (Neptunea) andersoni* (Martin) variety *hawleyi* (Carson)
Grant and Gale, 1931, p. 655, pl. 28, fig. 9a-9c.
Hypotype; 317; Pliocene, Pico Formation; Sulphur Canyon, Ventura Co., Calif.
- Ocenebra squamulifera* (Carpenter)
See: *T. [rophon] squamulifera* Carpenter
- Odostomia* cf. (*Amaura*) *avellana* Carpenter
Grant and Gale, 1931, p. 954, pl. 32, fig. 20.
Hypotype; 304; Pleistocene; southwest of Goleta, Santa Barbara Co., Calif.
- Odostomia* cf. (*Evalea*) *phanea* Dall and Bartsch
Grant and Gale, 1931, p. 933, pl. 24, fig. 25.
Hypotype; 218; Pleistocene; southwest of Goleta, Santa Barbara Co., Calif.
- Olivella biplicata* Sowerby
Grant and Gale, 1931, p. 625, pl. 24, fig. 15.
Hypotype; 211; Pleistocene; southwest of Goleta, Santa Barbara Co., Calif.
- Olivella ischnon* Keen
Keen, 1943, p. 50.
Paratype; 613, 937; Miocene, Round Mountain Silt or Olcese Sand; Caliente Quadrangle, Kern Co., Calif.
- Olivella pedroana* (Conrad)
Grant and Gale, 1931, p. 626, pl. 24, fig. 10.
Hypotype; 206; Pleistocene; southwest of Goleta, Santa Barbara Co., Calif.
- Olivella santana* Loel and Corey
Loel and Corey, 1932, p. 240.
Paratype; 795, 796; Miocene, Vaqueros Formation; north of Coal Mine on south side of 1332' hill, Santa Ana Mountains, Orange Co., Calif.
- Pseudomelatoma penicillata* (Carpenter) variety *semiinflata* Grant and Gale
Grant and Gale, 1931, p. 561, pl. 26, fig. 19.
Holotype; 237; Pleistocene; Santa Monica, Los Angeles Co., Calif.
- Puncturella punctostata* Berry
Berry, 1947, p. 265 (11).
Paratype; 325, 326; Pleistocene, Lomita Formation; near 2nd and Pacific Streets, San Pedro, Los Angeles Co., Calif.

Puncturella ralphi Berry

Berry, 1947, p. 267 (13).

Paratype; 327; Pleistocene; 2nd and Pacific Streets, San Pedro, Los Angeles Co., Calif.

Purpura (Jaton) eldridgei (Arnold)

Grant and Gale, 1931, p. 708, pl. 32, fig. 1, 2a, 2b.

Hypotype; 288 (fig. 1), 289 (fig. 2a, 2b); Pliocene, Pico Formation; east of Fernando Pass, Los Angeles Co., Calif.

Purpura (Jaton) festiva (Hinds)

Grant and Gale, 1931, p. 708, pl. 32, fig. 3.

Hypotype; 290; Pleistocene; San Pedro, Los Angeles Co., Calif.

Ranella (Priene) oregonensis (Redfield)

Grant and Gale, 1931, p. 737, pl. 27, fig. 12.

Hypotype; 269; Pliocene, Pico Formation; Sulphur Canyon, Los Angeles Co., Calif.

Rapana serrai Wiedey

Wiedey, 1928, p. 116, pl. 9, fig. 4-6.

Holotype; 13; Miocene, Vaqueros Formation; Kavannah Creek, San Luis Obispo Co., Calif.

Retusa (Acteocina) culicella (Gould)

Grant and Gale, 1931, p. 447, pl. 24, fig. 13.

Hypotype; 209; Pleistocene; Harmon Canyon, Ventura Co., Calif.

Scissurella lyra Berry

Berry, 1947, p. 268 (14).

Paratype; 328, 329; Pleistocene; 2nd and Pacific Streets, San Pedro, Los Angeles Co., Calif.

Searlesia branneri Clark and Arnold

(Clark and Arnold, 1923, p. 159, pl. 30, fig. 3a, 3b).

Plastoholotype; 22, 717; Oligocene or Miocene; Sooke Formation; Vancouver Island, British Columbia, Canada.

Solenosteira merriami Loel and Corey

Loel and Corey, 1932, p. 242.

Paratype; 797; Miocene, Vaqueros Formation; ridge between Vaqueros and Reliz creeks, Monterey Co., Calif.

Spirotropis (Antiplanes) cf. bulimoides (Dall)

Grant and Gale, 1931, p. 557, pl. 26, fig. 26.

Hypotype; 243; Pliocene, Pico Formation; Elsmere Canyon, Los Angeles Co., Calif.

Spirotropis (Antiplanes) perversa (Gabb)

Grant and Gale, 1931, p. 553, pl. 26, fig. 22, 23a, 23b.

Hypotype; 239 (fig. 22), 240 (fig. 23a, 23b); Pleistocene; San Pedro, Los Angeles Co., Calif.

Spirotropis (Antiplanes) perversa (Gabb) variety *fernandoensis* (English)

Grant and Gale, 1931, p. 557, pl. 26, fig. 25.

Hypotype; 242; Pliocene, Pico Formation; Elsmere Canyon, Los Angeles Co., Calif.

Spirotropis (Antiplanes) perversa (Gabb) variety *pedroana* (Arnold)

Grant and Gale, 1931, p. 556, pl. 26, fig. 24, 32.

Hypotype; 241 (fig. 24), 246 (fig. 32); Pleistocene; San Pedro, Los Angeles Co., Calif.

Spirotropis (Typhlomangelia) cf. renaudi (Arnold)

Grant and Gale, 1931, p. 549, pl. 26, fig. 29.

Hypotype; 244; Pliocene, Pico Formation; Sulphur Canyon, Ventura Co., Calif.

Strombina recurva (Sowerby)

Grant and Gale, 1931, p. 699, pl. 26, fig. 38, 41.

Hypotype; 249 (fig. 38), 252 (fig. 41); Pleistocene?; Albemarle Island, Galapagos Islands.

Surculites (Megasurcula) carpenterianus (Gabb)

Grant and Gale, 1931, p. 497, pl. 25, fig. 4a, 4b.

Hypotype; 228; Pleistocene; east of Sexton Canyon, Ventura Co., Calif.

Surculites (Megasurcula) remondii (Gabb)

Grant and Gale, 1931, p. 495, pl. 25, fig. 5, 6.

Hypotype; 229 (fig. 5), 230 (fig. 6); Pliocene, Pico Formation; east of Fernando Pass, Los Angeles Co., Calif.

Tegula (Chlorostoma) funebralis (A. Adams)

Grant and Gale, 1931, p. 826, pl. 32, fig. 28, 29a, 29b.

Hypotype; 309 (fig. 28), 310 (fig. 29a, 29b); Pliocene, Pico Formation; 309; between Santa Paula Creek and Timber Canyon; 310; Canada de Aliso; Ventura Co., Calif.

Tegula (Chlorostoma) gallina (Forbes)

Grant and Gale, 1931, p. 827, pl. 32, fig. 30, 31.

Hypotype; 311 (fig. 30), 312 (fig. 31); Pliocene, Pico Formation; west (311) and east (312) of Fernando Pass, Los Angeles Co., Calif.

Terebra (Strioterebrum) cf. dislocata (Say)

Grant and Gale, 1931, p. 468, pl. 24, fig. 19.

Hypotype; 214; Pleistocene; Ventura, Ventura Co., Calif.

- Terebra* (subgenus =?) *santana* Loel and Corey
 Loel and Corey, 1932, p. 236.
 Paratype; 790-793; Miocene; Vaqueros Formation; west side of Plano Trabuco, Orange Co., Calif.
- Thais* (*Nucella*) *elsinorensis* Grant and Gale
 Grant and Gale, 1931, p. 719, pl. 32, fig. 13.
 Holotype; 299; Pliocene, Pico Formation; east of Fernando Pass, Los Angeles Co., Calif.
- Thais* (*Nucella*) *shumanaensis* Carson
 Carson, 1926, p. 56.
 Paratype; 629; Pliocene, Fernando Formation; railway cut 1/2 mile north of Schuman, Santa Barbara Co., Calif.
- Tritonalia forcipata* (Hinds)
 Grant and Gale, 1931, p. 709, pl. 32, fig. 11.
 Hypotype; 297; Pleistocene; Adams Canyon, Ventura Co., Calif.
- Tritonalia lurida* (Middendorff) variety *aspera* (Baird)
 Grant and Gale, 1931, p. 711, pl. 32, fig. 4.
 Hypotype; 291; Pleistocene; southwest of Goleta, Santa Barbara Co., Calif.
- Tritonalia lurida* (Middendorff) variety *munda* (Carpenter)
 Grant and Gale, 1931, p. 712, pl. 32, fig. 5.
 Hypotype; 292; Pleistocene; southwest of Goleta, Santa Barbara Co., Calif.
- Tritonalia poulosi* (Nuttall in Carpenter)
 Grant and Gale, 1931, p. 712, pl. 32, fig. 10.
 Hypotype; 296; Pleistocene; San Pedro, Los Angeles Co., Calif.
- Tritonalia ynezana* Loel and Corey
 Loel and Corey, 1932, p. 248.
 Paratype; 798-800; Miocene, Vaqueros Formation; 2.5 miles southwest of Buellton, Santa Barbara Co., Calif.
- Trophon* (*Boreotrophon*) *orpheus* (Gould)
 Grant and Gale, 1931, p. 722, pl. 32, fig. 9.
 Hypotype; 295; Pliocene, Pico Formation; Sulphur Canyon, Ventura Co., Calif.
- T. [trophon] squamulifer* Carpenter
 Carpenter in Gabb, 1869, p. 44.
 Plastoholotype; 952a, 952b, 953; Pleistocene; Santa Barbara, Santa Barbara Co., Calif.
 See: *Ocenebra squamulifer* (Carpenter)
- Turbonilla* (*Pyrgiscus*) *bravoensis* Keen
 Keen, 1943, p. 51.
 Paratype; 614, 934; Miocene, Round Mountain Silt; Caliente Quadrangle, Kern Co., Calif.
- Turbanilla* (*Pyrgolampros*) *idae* (T. S. Oldroyd)
 Grant and Gale, 1931, p. 869, pl. 24, fig. 23.
 Hypotype; 316; Pleistocene; Kalorama Canyon, Ventura, Ventura Co., Calif.
- Turbanilla* (*Pyrgolampros*) *mariposa* Keen
 Keen, 1943, p. 52.
 Paratype; 615, 935; Miocene, Round Mountain Silt; Caliente Quadrangle, Kern Co., Calif.
- Turcicula* *santacruzana* Arnold
 Arnold, 1908, p. 373.
 Paratype; 625; Oligocene, San Lorenzo Formation; on San Lorenzo River 3 miles above town of Boulder Creek, Santa Cruz Co., Calif.
- Turritella* *bosci* Hertlein and Jordan
 Wiedey, 1928, p. 117, pl. 10, fig. 7, pl. 11, fig. 1, 2.
 Hypotype; 44 (fig. 7), 45 (fig. 1), 46 (fig. 2); Miocene, Temblor Formation; 44: Barker's Ranch, Kern Co., Calif.; 45, 46: 2 miles southeast of El Modena, Orange Co., Calif.
- Turritella* *cooperi* Carpenter
 Grant and Gale, 1931, p. 771, pl. 24, fig. 28-34.
 Hypotype; 221 (fig. 28), 222 (fig. 29), 223 (fig. 30), 224 (fig. 31), 225 (fig. 32), 226 (fig. 33), 227 (fig. 34); Pleistocene; San Pedro, Los Angeles Co., Calif.
- Turritella* *inezana* Conrad
 Wiedey, 1928, p. 120, pl. 12, fig. 2.
 Hypotype; 48; Miocene, Vaqueros Formation; 5 miles east of San Luis Obispo, San Luis Obispo Co., Calif.
- Turritella* *inezana* Conrad var. *pertunuda* Wiedey
 Wiedey, 1928, p. 119, pl. 12, fig. 1.
 Holotype; 14; Miocene, Vaqueros Formation; Canyon de Piedra, San Luis Obispo Co., Calif.
- Turritella* *inezana* Conrad var. *sespcensis* Arnold
 Wiedey, 1928, p. 121, pl. 12, fig. 4.
 Hypotype; 49; Miocene, Vaqueros Formation; Squaw Flat, Ventura Co., Calif.
- Turritella* *jewettii* Carpenter
 Grant and Gale, 1931, p. 770, pl. 24, fig. 26, 27.
 Hypotype; 219 (fig. 26), 220 (fig. 20); Pleistocene; San Pedro, Los Angeles Co., Calif.

Turritella montereyana Wiedey

Wiedey, 1928, p. 123, pl. 21, fig. 3.

Syntype; 51; Miocene, Monterey Formation; 1.5 miles south of San Antonio River, Monterey Co., Calif.

Turritella ocoyana Conrad

Wiedey, 1928, p. 120, pl. 10, fig. 1, 3.

Hypotype; 43 (fig. 3; missing), 47 (fig. 1); Miocene, Temblor Formation; 2 miles west of Simmler, San Luis Obispo Co., Calif.

Turritella temblorensis Wiedey

Wiedey, 1928, p. 122, pl. 11, fig. 4.

Holotype; 14; Miocene, Temblor Formation; Santa Monica Mtns., Los Angeles Co., Calif.

Turritella vanrecki Arnold

Grant and Gale, 1931, p. 773, pl. 24, fig. 22.

Hypotype; 217; Pliocene, Pico Formation; Holser Canyon, Los Angeles Co., Calif.

Turritella variata Conrad

Wiedey, 1928, p. 120, pl. 12, fig. 8.

Hypotype; 50; Miocene, Vaqueros Formation; 5 miles east of San Luis Obispo, San Luis Obispo Co., Calif.

Vasum (Vasum) pufferi Emerson

Emerson, 1964, p. 11.

Homeotype; 703, 747; Pliocene, Imperial Formation; Coyote Mountain (= Carrizo Mountain), Imperial Co., Calif.

Vexillum (Uromitra) healeyi Fargo

Fargo, 1948, p. 110.

Paratype; 909-913; Pliocene; north part of St. Petersburg, Pinellas Co., Florida.

Viviparus washingtonianus Arnold and Hannibal

Hannibal, 1913, p. 194.

Paratype; 626; Eocene, Tejon Formation; on Olequa Creek 2 miles north of Little Falls, Washington.

Volvula gluma Keen

Keen, 1943, p. 54.

Paratype; 616; Miocene, Round Mountain Silt; Caliente Quadrangle, Kern Co., Calif.

ARTHROPODA

Trilobita

Achatella carleyi (Meek)

Howell, 1951, p. 287, pl. 8, fig. 5.

Hypotype; 877; Ordovician?, Fairmount Formation?; Cincinnati, Ohio.

Anchiopsis anchiops (Green)

Howell, 1951, p. 286, pl. 8, fig. 3.

Plastoholotype; 875; Devonian; Ulster Co., New York.

See: *Calymene anchiops* Green

Bathyurus extans (Hall)

Howell, 1951, p. 264, pl. 1, fig. 5-7.

Hypotype; 828 (fig. 5), 829 (fig. 6), 830 (fig. 7); Ordovician, Black River Formation; Great Bend, Jefferson Co., New York.

Brachymetopus cuyahogae (Claypole)

Howell, 1951, p. 274, pl. 6, fig. 1.

Hypotype; 854; Mississippian, Cuyahoga Formation; Akron, Ohio.

Bumastus armatus (Hall)

Howell, 1951, p. 268, pl. 3, fig. 4-6.

Hypotype; 840 (fig. 4), 841 (fig. 5), 842 (fig. 6); Silurian, Racine Formation; Bridgeport, Illinois.

Bumastus chicagoensis (Weller)

Howell, 1951, p. 267, pl. 3, fig. 3.

Hypotype; 839; Silurian, Racine Formation; Bridgeport, Illinois.

Bumastus cuniculus (Hall)

Howell, 1951, p. 269, pl. 3, fig. 7, pl. 5, fig. 1.

Hypotype; 843 (fig. 7); 847 (fig. 1); Silurian, Racine Formation; Wauwatosa, Wisconsin.

Bumastus ? vogdesi Howell

Howell, 1951, p. 269, pl. 3, fig. 2.

Holotype; 838; Silurian; Dug Gap, Walker Co., Georgia.

Calymene anchiops Green

Green, 1832, p. 35.

Plastoholotype; 875; Devonian; Ulster Co., New York.

See: *Anchiopsis anchiops* (Green)

- Calymene clintoni* (Vanuxem)
 Vogdes, 1880, p. 178, fig. 3.
 Hypotype; 865; Taylor's Ridge near Catoosa Station, Catoosa Co., Georgia.
 See: *Calymene vogdesi* Foerste
 Vogdes, 1886, p. 5, fig. 3.
 Hypotype; 865; Taylor's Ridge near Catoosa Station, Catoosa Co., Georgia
- Calymene rostrata* Vogdes
 Vogdes, 1879, p. 477.
 Syntype; 868-872; Silurian, Clinton Group; Taylor's Ridge, Catoosa Station, Catoosa Co., Georgia.
 See: *Calymenella rostrata* (Vogdes)
 Vogdes, 1880, p. 176, text-fig. 1, 2.
 Syntype; 868 (fig. 1), 870 (fig. 2); Silurian, Clinton Group; Taylor's Ridge, Catoosa Station, Catoosa Co., Georgia.
 Vogdes, 1886, p. 2, text-fig. 1, 2.
 Syntype; 868 (fig. 1), 870 (fig. 2); Silurian, Clinton Group; Taylor's Ridge, Catoosa Station, Catoosa Co., Georgia.
- Calymene vogdesi* Foerste
 Foerste, 1887, p. 95.
 See: *Calymene clintoni* (Vanuxem)
 Howell, 1951, p. 280, pl. 7, fig. 5-7.
 Hypotype; 865 (fig. 5), 866 (fig. 6), 867 (fig. 7); Silurian; 865: Taylor's Ridge near Catoosa Station, Catoosa Co., Georgia; 866, 867: Dug Gap, Walker Co., Georgia.
- Calymenella rostrata* (Vogdes)
 Howell, 1951, p. 283, pl. 7, fig. 8-10.
 Syntype; 868 (fig. 8), 869 (fig. 9), 870 (fig. 10), 871, 872; Silurian, Clinton Group; Taylor's Ridge, Catoosa Station, Catoosa Co., Georgia.
 See: *Calymene rostrata* Vogdes
- Ceratocephala anchoralis* (Miller)
 Howell, 1951, p. 277, pl. 7, fig. 2.
 Hypotype; 861; Ordovician, Maysville Formation; Cincinnati region, Ohio.
- Ceratocephala longispina* (Mitchell)
 Howell, 1951, p. 305, pl. 13, fig. 1-2.
 Hypotype; 900 (fig. 1), 901 (fig. 2); Silurian, Bowning Series; Bowning Village, County Harden, New South Wales, Australia.
- Ceraurus pleurexanthemus* Green
 Howell, 1951, p. 284, pl. 8, fig. 1, 2.
 Hypotype; 873 (fig. 1), 874 (fig. 2); Ordovician; 873: Trenton Formation?, New York; 874: Trenton Formation, Trenton Falls, New York.
- Corycephalus dentatus* (Barrett)
 Howell, 1951, p. 286, pl. 8, fig. 4.
 Hypotype; 876; Devonian; Port Jervis, New York.
- Cyphaspis christyi* Hall
 Howell, 1951, p. 274, pl. 5, fig. 7, 8.
 Hypotype; 852; Silurian, Niagara Group; ?Waldron Formation; ?Waldron, ?Indiana.
- Cyphaspis yassensis* Etheridge and Mitchell
 Howell, 1951, p. 303, pl. 12, fig. 1-2.
 Hypotype; 895 (fig. 1), 896 (fig. 2); Silurian, Bowning Series; Belle Vale, New South Wales, Australia.
- Encrinurus americanus* Vogdes
 Vogdes, 1886, p. 1.
 Syntype; 862, 863; Plastosyntype; 864; Silurian, Clinton Group; Taylor's Ridge, west of Catoosa Station, Catoosa Co., Georgia.
 Howell, 1951, p. 278, pl. 7, fig. 3, 4.
 Syntype; 862 (fig. 3), 863; plastosyntype; 864 (cast from 863) (fig. 4); Silurian, Clinton Group; Taylor's Ridge, west of Catoosa Station, Catoosa Co., Georgia.
- Eophacops catoosaeensis* Howell
 Howell, 1951, p. 289, pl. 10, fig. 2-4.
 Holotype; 882 (fig. 2); paratype; 883 (fig. 3), 884 (fig. 4), 885; Silurian, Clinton Group; bank of Chicamauga River at the Ringgold end of the bridge on the road from Catoosa Station to Ringgold, Georgia.
- Griffithides bufo* Meek and Worthen
 Howell, 1951, p. 277, pl. 6, fig. 6; pl. 7, fig. 1.
 Hypotype; 859 (pl. 6, fig. 6); Plastohypotype; 860 (pl. 7, fig. 1); Mississippian, Keokuk Group; Crawfordsville, Indiana.
- Griffithides conwayensis* Wheeler
 (Wheeler, 1935, p. 52, pl. 6, fig. 4, 5.)
 Plastoholotype; 388; Pennsylvanian, Atoka Formation; Conway Co., Arkansas.
 See: *Griffithides ornata* Vogdes

- Griffithides nosoniensis* Wheeler
 (Wheeler, 1935, p. 51, pl. 6, fig. 6, 7.)
 Plastoholotype; 389; Permian, Nosoni Formation; ridge between Potter and Marble Creeks, Shasta Co., Calif.
- Griffithides ornata* Vogdes
 (Vogdes, 1895, p. 589, text-fig.)
 Plastoholotype; 388; Pennsylvanian, Atoka Formation; Conway Co., Arkansas.
 See: *Griffithides conwayensis* Wheeler
- Isotelus gigas* Dekay
 Howell, 1951, p. 265, pl. 2, fig. 3, pl. 3, fig. 1, pl. 4, fig. 1.
 Hypotype; 835 (pl. 2, fig. 3); 837 (pl. 3, fig. 1); 846 (pl. 4, fig. 1); Ordovician; 835: Trenton Formation, Herkimer Co., New York; 837: Kentucky; 846: "probably Kentucky or Cincinnati region".
- Lloydia parva* Howell
 Howell, 1951, p. 263, pl. 1, fig. 3, 4.
 Holotype; 825 (fig. 3), 826 (fig. 4), 827; Cambrian, Levis Conglomerate (Cambrian pebble in Ordovician conglomerate); Point Levis, Quebec, Canada.
- Loganellus macropleurus* Rasetti
 Howell, 1951, p. 263, pl. 1, fig. 2.
 Hypotype; 824; Cambrian, Levis Conglomerate (Cambrian pebble in Ordovician conglomerate); Point Levis, Quebec, Canada.
- Odontopleura jenkinsi* Etheridge and Mitchell
 Howell, 1951, p. 304, pl. 12, fig. 4.
 Hypotype; 898; Silurian, Bowning Series; "presumably" Bowning railway station yard, Bowning Village, County Harden, New South Wales, Australia.
- Odonotopleura rattei* Etheridge and Mitchell
 Howell, 1951, p. 304, pl. 12, fig. 3.
 Hypotype; 897; Silurian, Bowning Series; Bowning Village, County Harden, New South Wales, Australia.
- Ogygites canadensis* (Chapman)
 Howell, 1951, p. 265, pl. 1, fig. 8, pl. 2, fig. 1, 2.
 Hypotype; 831 (fig. 8), 833 (fig. 1), 834 (fig. 2); Ordovician, Collingwood Formation; Georgian Bay, Ontario, Canada.
- Phacops cacapona* Hall
 Howell, 1951, p. 288, pl. 9, fig. 2-4.
 Hypotype; 879; Devonian ?; Hamilton Group ?; Capon Springs, West Virginia.
- Phacops crosslei* Etheridge and Mitchell
 Howell, 1951, p. 307, pl. 13, fig. 3-4.
 Hypotype; 902 (fig. 3), 903 (fig. 4); Silurian, Bowning Series; Bowning Village, County Harden, New South Wales, Australia.
- Phacops hudsonica* Hall
 Howell, 1951, p. 288, pl. 9, fig. 5, pl. 10, fig. 1.
 Hypotype; 880; Devonian; western Tennessee.
- Phacops rana* (Green)
 Howell, 1951, p. 288, pl. 9, fig. 1.
 Hypotype; 878; Devonian, Hamilton Group; London, Ontario, Canada.
- Phacops serratus* Foerste
 Howell, 1951, p. 307, pl. 12, fig. 5.
 Hypotype; 899; Silurian, Bowning Series; near the railway station, Bowning, County Harden, New South Wales, Australia.
- Phillipsia insignis* Winchell
 Howell, 1951, p. 275, pl. 5, fig. 9.
 Hypotype; 853; Mississippian, ?Burlington Limestone; Curryville, Missouri.
- Phillipsia major* Shumard
 Howell, 1951, p. 275, pl. 6, fig. 2, 3.
 Hypotype; 855 (fig. 2), 856 (fig. 3); Pennsylvanian; Kansas City, Missouri.
- Phillipsia sampsoni* Vogdes
 Howell, 1951, p. 276, pl. 6, fig. 4.
 Hypotype; 857; Mississippian, Chouteau Formation; Banks, Pettis Co., Missouri.
- Phillipsia stevensoni* Meek
 Howell, 1951, p. 276, pl. 6, fig. 5.
 Hypotype; 858; Mississippian, Chester Group; Monongalia Co., West Virginia.
- Proetus*, Species undetermined
 Howell, 1951, p. 270, pl. 5, fig. 2.
 Hypotype; 848; Silurian; Taylor's Ridge near Catoosa Station, Catoosa Co., Georgia.
- Proetus bairdensis* Wheeler
 (Wheeler, 1935, p. 49, pl. 6, fig. 1-3).
 Plastoholotype; 390 (fig. 1, 3); plastoparatype; 339 (fig. 1, 2); Pennsylvanian?. Baird Formation;

- Redding Quadrangle, Shasta Co., Calif.
 Howell, 1951, p. 272, pl. 5, fig. 6.
 Hypotype; 851; Pennsylvanian?, Baird Formation; Redding Quadrangle, Shasta Co., Calif.
- Proctus crassimarginatus* (Hall)
 Howell, 1951, p. 271, pl. 5, fig. 3.
 Hypotype; 849; Devonian, Onondaga Formation; Kelly Island, Lake Erie.
- Proctus doris* Hall
 Hall, 1860, p. 112.
 Plastoptyype; 845; Carboniferous, Goniatite Limestone; Rockford, Indiana.
 Howell, 1951, p. 271, pl. 3, fig. 8, 9.
 Plastohypotype; 844 (fig. 8); plastoptyype; 845 (fig. 9); Carboniferous, Goniatite Limestone; Rockford, Indiana.
- Proctus haldemanii* Hall
 Howell, 1951, p. 272, pl. 5, fig. 4, 5.
 Hypotype; 850; Devonian, Hamilton Formation; Judd's Falls, on the road between Cherry Valley and Sharon Springs, New York.
- Proctus parvusculus* Hall
 Howell, 1951, p. 270, pl. 1, fig. 9, 10.
 Hypotype; 832; Ordovician, Corryville Formation; Cincinnati, Ohio.
- Proctus rattei* Etheridge and Mitchell
 Howell, 1951, p. 303, pl. 11, fig. 5.
 Hypotype; 894; Silurian, Bowning Series; Bowning, County Harden, New South Wales, Australia.
- Richardsonella unisulcata* Rasetti
 Howell, 1951, p. 263, pl. 1, fig. 1.
 Hypotype; 823; Cambrian, Levis Conglomerate (Cambrian pebble in Ordovician conglomerate); Point Levis, Quebec, Canada.
- Scutellum bowningense* (Etheridge and Mitchell)
 Howell, 1951, p. 301, pl. 10, fig. 7.
 Hypotype; 888; Silurian, Bowning Series; Bowning Creek, County Harden, New South Wales, Australia.
- Scutellum jenkinsi* (Etheridge and Mitchell)
 Howell, 1951, p. 302, pl. 10, fig. 8; pl. 11, fig. 1-3.
 Hypotype; 889 (fig. 8), 890 (fig. 1), 891 (fig. 2), 892 (fig. 3); Silurian, Bowning Series; Bowning Creek, County Harden, New South Wales, Australia.
- Scutellum longispinifex* (Mitchell)
 Howell, 1951, p. 302, pl. 11, fig. 4.
 Hypotype; 893; Silurian, Bowning Series; Bowning, County Harden, New South Wales, Australia.
- Scutellum partschi* (Barrande)
 Howell, 1951, p. 293, pl. 10, fig. 5, 6.
 Hypotype; 886 (fig. 5), 887 (fig. 6); Silurian; Lochkov, Bohemia.
- Vogdesia vigilans* (Meek and Worthen)
 Howell, 1951, p. 267, pl. 2, fig. 4.
 Hypotype; 836; Silurian, Maquoketa Formation; Fayette County, Iowa.
- Arachnida
- Calciro fisheri* Petrunkevitch
 Petrunkevitch, 1945, p. 323, pl. 1, right fig.; text-fig. 6.
 Paratype; 1137; Middle Cenozoic or younger; Bonner Quarry, north side of Black Mesa, about 10 miles southwest of Ashfork, Yavapai Co., Arizona.
- Crustacea
- Actinocythereis allisoni* Holden
 Holden, 1964, p. 418, text-fig. 22a, 22b.
 Holotype; 1004 (fig. 22a); paratype; 1005 (fig. 22b); Cretaceous, Rosario Formation; near Carlsbad, San Diego Co., Calif.
- Ampibycythereira iniqua* Holden
 Holden, 1964, p. 414, text-fig. 19a, 19c, 19e-19g.
 Holotype; 994 (fig. 19a); paratype; 995 (fig. 19c); 996 (fig. 19e, 19f); 997a (fig. 19g); 997b; Cretaceous, Rosario Formation; near Carlsbad, San Diego Co., Calif.
- Argilloccia constricta* Holden
 Holden, 1964, p. 403, text-fig. 9a-9c.
 Holotype; 971; Cretaceous Rosario Formation; near Carlsbad, San Diego Co., Calif.
- Bairdopilata cretacea* Holden
 Holden, 1964, p. 400, text-fig. 7a-7c, 7f-7i.
 Holotype; 965 (fig. 7a-7c); paratype; 966 (fig. 7f), 967 (fig. 7g), 968 (fig. 7h), 969 (fig. 7i); Cretaceous, Rosario Formation; near Carlsbad, San Diego Co., Calif.

Brachycythere darenensis Swain

Holden, 1964, p. 405, text-fig. 11a-11c, 11e.

Holotype; 973 (fig. 11a); 974 (fig. 11b); 975 (fig. 11c); 976 (fig. 11e); Cretaceous, Rosario Formation; near Carlsbad, San Diego Co., Calif.

Callianassa stephensi Rathbun

Rathbun, 1926, p. 122.

Paratype; 920-925; Pleistocene; Spanish Bight, San Diego, San Diego Co., Calif.

Cancer branneri Rathbun

Rathbun, 1926, p. 63.

Paratype; 915; Pleistocene; San Pedro, Los Angeles Co., Calif.

Cythereis brooksi Holden

Holden, 1964, p. 419, text-fig. 23a-23c.

Holotype; 1006 (fig. 23a, 23b); paratype; 1007 (fig. 23c); Cretaceous, Rosario Formation; near Carlsbad, San Diego Co., Calif.

Cytherella elliotti Holden

Holden, 1964, p. 397, text-fig. 4b, 4c.

Holotype; 958; Cretaceous, Rosario Formation; near Carlsbad, San Diego Co., Calif.

Cytherella terminopunctata Holden

Holden, 1964, p. 396, text-fig. 3a, 3c, 3d.

Holotype; 955 (fig. 3c); paratype; 956 (fig. 3a); 957 (fig. 3d); Cretaceous, Rosario Formation; near Carlsbad, San Diego Co., Calif.

Cytherelloidea directiangula Holden

Holden, 1964, p. 399, text-fig. 6a-6e.

Holotype; 963 (fig. 6a-6c); paratype; 964 (fig. 6d, 6e); Cretaceous, Rosario Formation; near Carlsbad, San Diego Co., Calif.

Cytherelloidea milowi Holden

Holden, 1964, p. 398, text-fig. 5a-5e.

Holotype; 959 (fig. 5a); paratype; 960 (fig. 5b); 961 (fig. 5c); 962a (fig. 5d); 962b (fig. 5e); Cretaceous, Rosario Formation; near Carlsbad, San Diego Co., Calif.

Cytheropteron coryelli carlsbadensis Holden

Holden, 1964, p. 409, text-fig. 14a-14e.

Holotype; 981 (fig. 14a, 14c, 14d); paratype; 982 (fig. 14b); 983 (fig. 14e); Cretaceous, Rosario Formation; near Carlsbad, San Diego Co., Calif.

Cytherura (?) divaricata Holden

Holden, 1964, p. 407, text-fig. 13a, 13b, 13e.

Holotype; 978 (fig. 13a); paratype; 979 (fig. 13b); 980 (fig. 13e); Cretaceous, Rosario Formation; near Carlsbad, San Diego Co., Calif.

Eocytheropteron turgidulum Holden

Holden, 1964, p. 410, text-fig. 15a-15c.

Holotype; 984; Cretaceous, Rosario Formation; near Carlsbad, San Diego Co., Calif.

Eucytherura planolata Holden

Holden, 1964, p. 412, text-fig. 17a-17c.

Holotype; 988 (fig. 17a, 17b); paratype; 989 (fig. 17c); Cretaceous, Rosario Formation; near Carlsbad, San Diego Co., Calif.

Eucytherura spinata Holden

Holden, 1964, p. 413, text-fig. 18a-18e.

Holotype; 990 (fig. 18a, 18b); paratype; 991 (fig. 18c); 992 (fig. 18d); 993 (fig. 18e); Cretaceous, Rosario Formation; near Carlsbad, San Diego Co., Calif.

Eucytherura versabilis Holden

Holden, 1964, p. 410, text-fig. 16b-16e.

Holotype; 985 (fig. 16b); paratype; 986 (fig. 16c); 987 (fig. 16d, 16e); Cretaceous, Rosario Formation; near Carlsbad, San Diego Co., Calif.

Idiocythere triebeli Holden

Holden, 1964, p. 422, text-fig. 25a-25c.

Holotype; 1012 (fig. 25a, 25b); paratype; 1013 (fig. 25c); Cretaceous, Rosario Formation; near Carlsbad, San Diego Co., Calif.

Isocythereis carlsbadensis Holden

Holden, 1964, p. 423, text-fig. 26a-26f.

Holotype; 1014 (fig. 26a-26c); paratype; 1015 (fig. 26d-26f); 1016; Cretaceous, Rosario Formation; near Carlsbad, San Diego Co., Calif.

Krithe cushmani carlsbadensis Holden

Holden, 1964, p. 406, text-fig. 12a-12c.

Holotype; 977; Cretaceous, Rosario Formation; near Carlsbad, San Diego Co., Calif.

Neocythere (Physocythere) formicata Holden

Holden, 1964, p. 404, text-fig. 10a-10e.

Holotype; 972 (fig. 10a-10c); paratype; 1066 (fig. 10d); 1067 (fig. 10e); Cretaceous, Rosario Formation; near Carlsbad, San Diego Co., Calif.

Paijenborchella pseudotrigona Holden

Holden, 1964, p. 415, text-fig. 20a, 20b, 20d, 20e.

Holotype; 998 (fig. 20a, 20b); paratype; 999 (fig. 20d); 1000 (fig. 20e); Cretaceous, Rosario Formation; near Carlsbad, San Diego Co., Calif.

Paracypris fragilis Holden

Holden, 1964, p. 402, text-fig. 8a.

Holotype; 970; Cretaceous, Rosario Formation; near Carlsbad, San Diego Co., Calif.

Platycosta oena Holden

Holden, 1964, p. 421, text-fig. 24d-24g.

Holotype; 1008 (fig. 24g); paratype; 1009; 1010 (fig. 24d); 1011 (fig. 24e, 24f); Cretaceous, Rosario Formation; near Carlsbad, San Diego Co., Calif.

Randallia pleistocenica Rathbun

Rathbun, 1926, p. 77.

Paratype; 916.919; Pleistocene; San Pedro, Los Angeles Co., Calif.

Trachyleberis acuminata Holden

Holden, 1964, p. 416, text-fig. 21a-21d.

Holotype; 1001 (fig. 21a); paratype; 1002 (fig. 21b); 1003 (fig. 21c, 21d); Cretaceous, Rosario Formation; near Carlsbad, San Diego Co., Calif.

Unidentified ostracod

Holden, 1964, p. 426, text-fig. 28a-28b.

Hypotype; 1023 (fig. 28a); 1024 (fig. 28b); Cretaceous, Rosario Formation; near Carlsbad, San Diego Co., Calif.

Xestoleberis minuta Holden

Holden, 1964, p. 425, text-fig. 27a-27c.

Holotype; 1017a (fig. 27a, 27c); 1017b (fig. 27b); paratype; 1018-1022; Cretaceous, Rosario Formation; near Carlsbad, San Diego Co., Calif.

ECHINODERMATA

Cidaris sp.

Grant and Hertlein, 1938, p. 7.

Hypotype; 351; Pliocene, San Diego Formation; Pacific Beach, San Diego, San Diego Co., Calif.

See: *Hesperocidaris perplexa* H. L. Clark

Dendraster casseli Grant and Hertlein

Grant and Hertlein, 1938, p. 81.

Paratype; 30; Pliocene; about 6 miles slightly north of west of Newhall, Los Angeles Co., Calif.

Emerson and Hertlein, 1964, p. 365, fig. 5a-5c.

Hypotype; 814 (5a), 815 (5b), 816 (5c); Pliocene; Isla San José, Baja California, Mexico.

Dendraster cf. D. granti Durham

Emerson and Hertlein, 1964, p. 365, fig. 5f.

Hypotype; 817; Pliocene; Isla San José, Baja California, Mexico.

Dendraster vizcainensis Grant and Hertlein

Grant and Hertlein, 1938, p. 90, pl. 8, fig. 1-3.

Holotype; 881; Pleistocene; Punta Santa Rosalia, Bahía de Sebastian Vizcaino, Baja California, Mexico.

Encope californica Verrill

Emerson and Hertlein, 1964, p. 365, fig. 6a-6e.

Hypotype; 819 (6a-6c); 820 (6d-6e); Pleistocene; Isla Monserrate, Baja California, Mexico.

Encope grandis L. Agassiz

Emerson and Hertlein, 1964, fig. 6f-6i.

Hypotype; 821 (6f), 822 (6g-6i); Pleistocene; Isla Monserrate, Baja California, Mexico.

Encope tenuis Kew

Hertlein and Grant, 1960, p. 126, pl. 25, fig. 4.

Hypotype; 352; Pliocene, San Diego Formation; 31st St. and Logan Ave., San Diego, San Diego Co., Calif.

Hesperocidaris perplexa H. L. Clark

Hertlein and Grant, 1960, p. 105, pl. 24, fig. 12.

Hypotype; 351; Pliocene, San Diego Formation; Pacific Beach, San Diego, San Diego Co., Calif.

See: *Cidaris* sp.

Megapetalus lovenioides Clark

(Clark, 1929, p. 260, pl. 31, fig. 1-6).

Plastoholotype; 72, 730, 31; Miocene; divide between Sulphur and Coche canyons, Ventura Co., Calif.

Merriamaster cf. M. israelskyi E. K. Jordan and Hertlein

Hertlein and Grant, 1960, p. 122, pl. 23, fig. 10.

Hypotype; 350; Pliocene, San Diego Formation; Cabrillo Canyon, Balboa Park, San Diego, San Diego Co., Calif.

- Oligopygus putnami* Israelsky
 (Israelsky, 1933, p. 275, pl. 18, fig. 1, 2, 4).
 Plastoptype; 422 (fig. 1), 423 (fig. 2), 424 (fig. 4); Eocene; 12 kilometers northeast of Abasola, Tamaulipas, Mexico.
- Scutaster vaquerosensis* var. *kewi* Loel and Corey
 Loel and Corey, 1932, p. 180.
 Paratype; 774; Miocene, Vaqueros Formation; west side of Grimes Canyon, about one mile from the mouth, Ventura Co., Calif.
- Spatangus tapinus* Schenck
 Schenck, 1928, p. 198, pl. 24, fig. 1, 3, 4.
 Holotype; 52; Eocene, Tejon Formation; Timber Canyon, Ventura Co., Calif.

LITERATURE CITED

- ARNOLD, R.
- 1903. The paleontology and stratigraphy of the marine Pliocene and Pleistocene of San Pedro, California. Mem. Calif. Acad. Sci., Vol. 3. 420 pp., 37 pl.
 - 1908. Descriptions of new Cretaceous and Tertiary fossils from the Santa Cruz Mountains, California. Proc. U.S. Nat. Mus. 34:345-389, pl. 31-36.
- BELL, W. A.
- 1962. Catalogue of types and figured specimens of fossil plants in the Geological Survey of Canada collections. Dept. Mines and Tech. Surveys of Canada. Ottawa. viii + 154 pp.
- BERRY, S. S.
- 1940. New Mollusca from the Pleistocene of San Pedro, California-I. Bull. Amer. Paleont. 25 (94A):149-164 (1-18), pl. 17-18 (1-2).
 - 1941. New Mollusca from the Pleistocene of San Pedro, California-II. Bull. Amer. Paleont. 27(101):1-18, pl. 1.
 - 1947. New Mollusca from the Pleistocene of San Pedro, California-III. Bull. Amer. Paleont. 31(127):257-274 (1-20), pl. 26-27 (1-2).
 - 1954. New Californian Pleistocene Eulimidae. Bull. Amer. Paleont. 35(151):257-270 (1-16), pl. 24 (1).
- CARSON, C. M.
- 1926. New molluscan species from the Californian Pliocene. Bull. So. Calif. Acad. Sci. 25:49-62.
- CLARK, B. L.
- 1925. Pelecypoda from the marine Oligocene of western North America. Univ. Calif. Publ. Geol. Sci. 15:69-136.
- CLARK, B. L., AND R. ARNOLD
- 1923. Fauna of the Sooke Formation, Vancouver Island. Univ. Calif. Publ. Geol. Sci. 14:123-234.
- CLARK, H. L.
- 1929. A new Miocene echinoid from California. Trans. San Diego Soc. Nat. Hist. 5:257-262, pl. 31.
- CUSHMAN, J. A., AND U. S. GRANT, IV.
- 1927. Late Tertiary and Quaternary Elphidiums of the west coast of North America. Trans. San Diego Soc. Nat. Hist. 5:69-82, pl. 7, 8.
- CUSHMAN, J. A., AND M. A. HANNA
- 1927. Foraminifera from the Eocene near San Diego, California. Trans. San Diego Soc. Nat. Hist. 5:45-64, pl. 4-6.
- CUSHMAN, J. A., R. E. STEWART, AND K. C. STEWART
- 1930. Tertiary foraminifera from Humboldt County, California. Trans. San Diego Soc. Nat. Hist. 6:41-94, pl. 1-8, 1 chart.
- DALL, W. H.
- 1921. New fossil invertebrates from San Quentin Bay, Lower California. West Amer. Sci. 19:17-18.

- DAVIS, C. H.
- 1913. New species from the Santa Lucia Mountains, California, with a discussion of the Jurassic age of the slates at Slate's Springs. Jour. Geol. 21:453-458.
- EMERSON, W. K.
- 1964. Results of the Puritan-American Museum of Natural History Expedition to western Mexico. 20. The Recent Mollusks: Gastropoda: Harpidae, Vasidae, and Volutidae. Amer. Mus. Novitates no. 2202. 23 pp.
- EMERSON, W. K., AND L. G. HERTLEIN
- 1964. Invertebrate megafossils of the Belvedere expedition to the Gulf of California. Trans. San Diego Soc. Nat. Hist. 13:333-368.
- FARGO, W. G.
- 1948. New *Vexillum* and *Aesopus* from the Pliocene of St. Petersburg, Florida. Nautilus 61:109-112, pl. 7.
- FOERSTE, A. F.
- 1887. The Clinton group of Ohio. Part II. Bull. Sci. Lab. Denison Univ. 2:89-110, pl. 8.
- FRIZZELL, D. L.
- 1930. A new Pleistocene fossil from Port Blakely, Washington. Nautilus 43:120-121.
 - 1931. A molluscan species new to the Recent west coast fauna. Trans. San Diego Soc. Nat. Hist. 6:319-324, pl. 22.
- GABB, W. M.
- 1869. Palaeontology of California. Vol. 2. Cretaceous and Tertiary fossils. Geol. Survey Calif. 229 pp., pl. 1-36.
- GRANT, U. S., IV, AND H. R. GALE
- 1931. Catalogue of the Marine Pliocene and Pleistocene Mollusca of California and Adjacent Regions. Mem. San Diego Soc. Nat. Hist., Vol. 1. 1036 pp.
- GRANT, U. S., IV, AND L. G. HERTLEIN
- 1938. The west American Cenozoic Echinoidea. Publ. Univ. Calif. Los Angeles Math. Phys. Sci., Vol. 2. 226 pp.
- GREEN, J.
- 1832. A Monograph of the Trilobites of North America: with Coloured Models of the Species. Joseph Brano, Philadelphia. 94 pp., 1 pl.
- HALL, J.
- 1860. Notes and observations upon the fossils of the Goniatite limestone in the Marcellus shale of the Hamilton group, in the eastern and central parts of the state of New York, and those of the Goniatite beds of Rockford, Indiana; with some analogous forms from the Hamilton group proper. Regents on the [N. Y.] State Cabinet, 13th Ann. Rep., pp. 95-112.
- HANNA, G. D.
- 1926. Paleontology of Coyote Mountain, Imperial County, California. Proc. Calif. Acad. Sci., ser. 4, 14:427-503.
- HANNIBAL, H.
- 1913. A synopsis of the Recent and Tertiary freshwater Mollusca of the Californian Province, based upon an ontogenetic classification. Proc. Malacol. Soc. London 10:112-211, pl. 5-8.
- HERTLEIN, L. G.
- 1925a. Pectens from the Tertiary of Lower California. Proc. Calif. Acad. Sci., ser. 4, 14:1-35.
 - 1925b. New species of marine fossil Mollusca from western North America. Bull So. Calif. Acad. Sci. 24:39-46.
 - 1933. A new gryphaeoid oyster from the Eocene of California. Trans. San Diego Soc. Nat. Hist. 7:277-279, pl. 18, figs. 5, 6.
 - 1934. New oysters and a new *Pecten* from the Tertiary of California. Bull. So. Calif. Acad. Sci. 33:1-6.

HERTLEIN, L. G., AND U.S. GRANT, IV

1944. The Cenozoic Brachiopoda of western North America. Univ. Calif. Los Angeles Publ. Math. Phys. Sci. Vol. 3. 236 pp.
1960. The geology and paleontology of the marine Pliocene of San Diego, California. Mem. San Diego Soc. Nat. Hist. Vol. 2, part 2a, pp. 73-133, pl. 19-26.

HOLDEN, J. C.

1964. Upper Cretaceous ostracods from California. Palaeont. 7:393-429.

HOWELL, B. F.

1951. The Vogdes collection of trilobites. Trans. San Diego Soc. Nat. Hist. 11:257-328, pl. 1-13.

INTERNATIONAL COMMISSION ON ZOOLOGICAL NOMENCLATURE

1964. International code of zoological nomenclature, adopted by the XV International Congress of Zoology. [N. R. STOLL (Chairman, Editorial Committee)] London. [International Trust for Zoological Nomenclature] xix + 176 pp.

ISRAELSKY, M. C.

1933. A new species of echinoid from Tamaulipas, Mexico. Trans. San Diego Soc. Nat. Hist. 7:275-276, pl. 18, figs. 1-4.

KANAKOFF, G. P.

1956. Two new species of *Nassarius* from the Pliocene of Los Angeles County, California. Bull. So. Calif. Acad. Sci. 55:110-114.

KEEN, A. M.

1943. New mollusks from the Round Mountain Silt (Tremblor) Miocene of California. Trans. San Diego Soc. Nat. Hist. 10:25-60, pl. 3-4.

LAMY, E.

1921. Revision des Carditacea vivants du Muséum National d'Histoire Naturelle de Paris. Jour. de Conch. 66:218-276.

LOEL, W., AND W. H. COREY

1932. The Vaqueros Formation, Lower Miocene of California. I. Paleontology. Univ. Calif. Publ. Geol. Sci. 22:31-140, 2 maps.

MEEK, F. B.

1857. Descriptions of new organic remains from the Cretaceous rocks of Vancouver's Island. Trans. Albany Inst. 4:37-49.

NICOL, D.

1944. New west American species of the foraminiferal genus *Elphidium*. Jour. Paleont. 18:172-185, pl. 29.

NOMLAND, J. O.

1916. Corals from the Cretaceous and Tertiary of California and Oregon. Univ. Calif. Publ. Geol. 9:59-76.

OLDROYD, T. S.

1921. New Pleistocene mollusks from California. Nautilus 34:114-116, pl. 5, fig. 8-13.

PETRUNKEVITCH, A.

1945. *Calcitra fisheri*. A new fossil arachnid. Amer. Jour. Sci. 243:320-329, pl. 1.

PILSBRY, H. A., AND A. A. OLSSON

1935. New mollusks from the Panamic Province. Nautilus 49:16-19, pl. 1.

QUAYLE, E. H.

1932. Fossil corals of the genus *Turbinolia* from the Eocene of California. Trans. San Diego Soc. Nat. Hist. 7:91-110, pl. 6.

RATHBUN, M. J.

1926. The fossil stalk-eyed Crustacea of the Pacific slope of North America. U. S. Nat. Mus. Bull. 138. vii + 155 pp., 39 pl.

- REINHART, P. W.
1937. Cretaceous and Tertiary pelecypods of the Pacific slope incorrectly assigned to the family Arcidae. *Jour. Paleont.* 11:169-180, pl. 27.
- RIVERS, J. J.
1913. A new species of *Bathytoma* from the Upper Pleistocene of San Pedro, Cal. *Bull. So. Calif. Acad. Sci.* 12:29, 1 pl.
- SCHENCK, H. G.
1928. A new echinoid from the California Eocene. *Trans. San Diego Soc. Nat. Hist.* 5:195-202, pl. 24.
1929. *Discocyclina* in California. *Trans. San Diego Soc. Nat. Hist.* 5:211-240, pl. 27-30.
- SCHENK, E. T., H. H. McMASTERS, A. M. KEEN, AND S. W. MULLER
1956. Procedure in Taxonomy. Stanford University Press, Stanford, Calif. vii + 149 pp.
- TEGLAND, N. M.
1931. The gastropod genus *Galeodea* in the Oligocene of Washington. *Univ. Calif. Publ. Geol. Sci.* 19:397-434.
1933. The fauna of the type Blakeley Upper Oligocene of Washington. *Univ. Calif. Publ. Geol. Sci.* 23:81-174.
- VOGDES, A. W.
1879. Short notes upon the geology of Catoosa County, Georgia. *Amer. Jour. Sci. and Arts*, ser. 3, 18:475-477.
1880. Description of new crustacean from the Upper Silurian of Georgia, with remarks upon *Calymene clintoni*. *Proc. Acad. Nat. Sci. Phila.* 1800:176-178.
1886. Description of new crustacean from the Clinton Group of Georgia, with remarks upon others. New York City. 5 pp. (privately published)
1895. Notes on Palaeozoic Crustacea no. 4 — On a new trilobite from Arkansas Lower Coal Measures. *Proc. Calif. Acad. Sci.*, ser. 2, 4:589-591.
- WARING, C. A.
1917. Stratigraphic and faunal relations of the Martinez to the Chico and Tejon of southern California. *Proc. Calif. Acad. Sci.*, ser. 4, 7:41-124.
- WHEELER, H. E.
1935. New trilobite species from the Anthracolithic of northern California and *Griffithides conwayensis*, a new name for a trilobite species from the Atoka Formation of Arkansas. *Trans. San Diego Soc. Nat. Hist.* 8:47-56, pl. 6.
- WIEDEY, L. W.
1928. Notes on the Vaqueros and Temblor formations of the California Miocene with descriptions of new species. *Trans. San Diego Soc. Nat. Hist.* 5:95-182, pl. 9-21.
1929. New Miocene mollusks from California. *Jour. Paleont.* 3:280-289, pl. 31-33.
- WOODRING, W. P.
1930. Upper Eocene orbitoid foraminifera from the western Santa Ynez Range, California, and their stratigraphic significance. *Trans. San Diego Soc. Nat. Hist.* 6:145-170, pl. 13-17.

SPECIES INDEX

- abbotti, *Spisula*, 109
 abietis, *Chlamys* (*Argopecten*), 103
 acolasta, *Macoma moesta*, 104
 acuminata, *Trachyleberis*, 122
 acutilineata, *Lucina* (*Myrtea*), 104
 albaria, *Mactra* (*Spisula*), 105
 alexi, *Eohemithiris*, 101
 allisoni, *Actinocythereis*, 120
 americanus, *Encriurus*, 118
 anchiops, *Anchiopsis*, 117
 anchiops, *Calymene*, 117
 anchoralis, *Ceratocephala*, 118
 anchuela, *Mitrella* (*Mitrella*), 113
 andersoni, *Neptunea* (*Neptunea*), 114
 andersoni, *Pecten* (*Aequipecten*), 106
 angelica, *Ostrea*, 105
 antillarum, *Crassatellites*, 103
 approximata, *Lucina* (*Myrtea*) *tenuisculpta*, 104
 apta, *Galeoidea*, 112
 arcumbona, *Cardium*, 103
 arenaria, *Mya* (*Mya*), 105
 aresta, *Margarites* (*Lirularia*), 113
 armatus, *Bumastus*, 117
 arnoldi, *Nassarius* (*Uzita*), 114
 ashleyi, *Ostrea*, 105
 aspera, *Tritonalia lurida*, 116
 asperum, *Bittium* (*Lirobittium*), 110
 aster, *Actinocyclina*, 99
 avellana, *Odostomia* (*Amaura*), 114
 bairdensis, *Proetus*, 119
 bakeri, *Pecten* (*Janira*) *stearnsii*, 107
 balthica, *Macoma*, 104
 barbarens, *Elphidium fax*, 100
 barbarensis, *Fusinus*, 112
 bellus, *Pecten* (*Janira*), 107
 beringianus, *Pecten* (*Pecten*), 107
 biplicata, *Olivella*, 114
 birchi, *Nucula* (*Ennucula*), 105
 bisecta, *Thyasira*, 109
 bodegensis, *Tellina*, 109
 bosei, *Turritella*, 116
 bowningense, *Scutellum*, 120
 branneri, *Cancer*, 121
 branneri, *Searlesia*, 115
 braeensis, *Turbonilla* (*Pyrgiscus*), 116
 brooksi, *Cythereis*, 121
 bufo, *Griffithides*, 118
 bulimoides, *Spirotropis* (*Antiplanes*), 115
 cacapona, *Phacops*, 119
 californianus, *Nassarius* (*Schizopyga*), 113
 californianus, *Nassarius* (*Tritia*), 114
 californianus, *Tagelus*, 109
 californica, *Amphistegina*, 99
 californica, *Cerithidea*, 111
 californica, *Cryptomya*, 103
 californica, *Discocyclina*, 99
 californica, *Encope*, 122
 californica, *Lucina* (*Myrtea*), 104
 californicus, *Conus*, 111
 callidus, *Pecten* (*Aequipecten*) *purpuratus*, 106
 callosa, *Amiantis*, 101
 camuloensis, *Arca* (*Arca*) *multicostata*, 102
 canadensis, *Ogygites*, 119
 canaliculatum, *Calliostoma*, 110
 cancellata, *Mya* (*Platyodon*), 105
 carditoides, *Petricola*, 108
 careyi, *Serpula*, 101
 carinata, *Mitrella*, 113
 carleyi, *Achatella*, 117
 carlsbadensis, *Cytheropteron coryelli*, 121
 carlsbadensis, *Isocythereis*, 121
 carlsbadensis, *Kritte cushmani*, 121
 carpenteri, *Cardita* (*Carditamera*), 103
 carpenterianus, *Surculites* (*Megasurcula*), 115
 carrizoensis, *Antigona*, 102
 casseli, *Dendraster*, 122
 catalinae, *Mitra*, 113
 catilliformis, *Mactra* (*Spisula*), 105
 catoosaensis, *Eophacops*, 118
 caurinus, *Pecten* (*Patinopecten*), 107
 chicagoensis, *Bumastus*, 117
 centrifilosum, *Laevicardium* (*Nemocardium*), 104
 cerrosensis, *Pecten* (*Lyropecten*) *estrellanus*, 107
 chacei, *Monilopsis*, 113
 christyi, *Cyphaspis*, 118
 clarki, *Turbinolia*, 101
 clarkiana, *Bathyoma*, 110
 clavatula, *Cancellaria*, 111
 clavella, *Balcis* (*Balcis*), 110
 clintoni, *Calymene*, 118
 coalingensis, *Pecten* (*Janira*) *bellus*, 107
 coalingensis, *Clavus* (*Clathrodrillia*), 111
 columbiana, *Amphissa*, 110
 communis, *Nodosaria* (*Dentalina*), 100
 compressa, *Architeconica*, 110
 cenchita, *Balcis*, 110
 conradiana, *Clathurella*, 111
 consobrina, *Nodosaria* (*Dentalina*), 100
 constricta, *Argilloecia*, 120
 contignata, *Ficus* (*Trophosycon*) *ocoyana*, 112
 convexa, *Gaudryina*, 100
 conwayensis, *Griffithides*, 118
 cooperi, *Nassarius* (*Schizopyga*) *mendicus*, 114
 cooperi, *Nassarius* (*Tritia*) *mendicus*, 114
 cooperi, *Turritella*, 116
 copelandi, *Macoma*, 104
 corbis, *Laevicardium* (*Cerastoderma*), 104
 coreyi, *Serpula*, 101
 coryelli, *Cytheropteron*, 121
 cretacea, *Bairdopplata*, 120
 cristobalensis, *Pecten* (*Plagioctenium*), 107
 crassimarginatus, *Proetus*, 120
 crosslei, *Phacops*, 119
 culcitella, *Retusa* (*Acteocina*), 115
 cuniculus, *Bumastus*, 117
 cushmani, *Kritte*, 121
 cuyahogae, *Brachymetopus*, 117
 cylcia, *Adontorhina*, 101
 darenensis, *Brachycythere*, 121
 decis, *Semele*, 108
 dentatus, *Corycehalus*, 118
 deserti, *Pecten* (*Aequipecten*), 106
 dicksoni, *Turbinolia*, 101
 diegensis, *Laqueus vancouverensis*, 101
 diegensis, *Pecten* (*Janira*) *stearnsii*, 107
 dilatatum, *Bittium* (*Lirobittium*) *asperum*, 110

- directiangula, Cytherelloidea, 121
 discus, Architectonica nobilis, 110
 discus, Pecten (Aequipecten), 106
 dislocata, Terebra (Strioterebrum), 115
 divaricata, Cytherura, 121
 divaricata, Lacuna, 112
 doris, Proetus, 120
 ebricanus, Balcis (Vitreolina), 110
 eldridgei, Ostrea, 105
 eldridgei, Purpura (Jaton), 115
 elliotti, Cytherella, 121
 elongata, Clementia, 103
 elsmensis, Thais (Nucella), 116
 elsmensis, Venus (Chione), 109
 eocenica, Epistomina, 100
 erici, Pecten (Chlamys), 106
 estrellanus, Pecten (Lyropecten), 107
 etchegoini, Pecten (Pallium) swiftii, 107
 excavata, Lucina (Here), 104
 extans, Bathyurus, 117
 fax, Elphidium, 100
 fernandoensis, Cancellaria tritonidea, 111
 fernandoensis, Pecten (Pseudamusium)
 vancouverensis, 108
 fernandoensis, Spirotropis (Antiplanes)
 perversa, 115
 fernandoensis, Venus (Chione) securis, 109
 festiva, Purpura (Jaton), 115
 fisheri, Calcitra, 120
 fornicata, Neocythere (Physocythere), 121
 fortis, Cantharus, 111
 fossatus, Nassarius (Schizopyga), 113
 fossatus, Nassarius (Tritia), 114
 fossilis, Conus californicus, 111
 foveolata, Tritonalia, 116
 fragilis, Paracypris, 122
 funebialis, Tegula (Chlorostoma), 115
 gabbi, Pholas (Zirfaea), 108
 galeana, Mitromorpha, 113
 galei, Arca, 102
 gallina, Tegula (Chlorostoma), 115
 gastoensis, Tivela, 109
 gausapata, Mitrella carinata, 113
 gemmulatum, Calliostoma, 111
 generosa, Panope (Panope), 106
 gettysburgensis, Nucula (Acila), 105
 gibbiformis, Corbula (Corbula), 103
 giganteus, Saxidomus nuttalli, 108
 gigas, Isotelus, 119
 gluma, Volvulella, 117
 gnomon, Hastula, 112
 gouldii, Donax, 103
 gracirosana, Moniliopsis, 113
 gradata, Astraea (Pomaularax), 110
 grandior, Acteon (Rictaxis) painei, 110
 grandior, Mitrella, 113
 grandis, Enope, 122
 granti, Dendraster, 122
 granti, Spisula, 109
 hakei, Pecten (Plagioctenium), 107
 haldemani, Proetus, 120
 haleyi, Ostrea, 105
 hamelini, Arca, 102
 hannibali, Dendrophyllia, 101
 harfordus, Pecten (Camptonectes), 106
 hawleyi, Neptunea (Neptunea) andersoni, 114
 healeyi, Pecten (Patinopecten), 107
 healeyi, Vexillum (Uromitra), 117
 hemphilli, Cancellaria, 111
 hemphilli, Pecten (Janira) bellus, 107
 hertleini, Basterotria (Basterotella), 103
 hildegardae, Nassarius, 113
 hodgei, Pecten (Chlamys), 106
 howelli, Ostrea, 105
 hudsonica, Phacops, 119
 hughesi, Elphidium, 100
 humerosus, Cantharus, 111
 idae, Tellina, 109
 idae, Turbonilla (Pyrgolampros), 116
 imbulata, Turbinolia, 101
 impavida, Arca, 102
 inaequalis, Astraea (Pachypoma), 110
 incallida, Balcis (Vitreolina), 110
 incisa, Moniliopsis incisa, 113
 inequalis, Clementia, 103
 inezana, Spondylus, 109
 inezana, Turritella, 116
 iniqua, Amphicytherura, 120
 inornata, Triloculina, 100
 inornatus, Robulus, 100
 insignis, Phillipsia, 119
 instabilis, Acmaea, 110
 invalidus, Pecten (Aequipecten) deserti, 106
 ischnon, Olivella, 114
 islandicus, Pecten (Pecten), 107
 israelskyi, Merriamaster, 122
 jacksonensis, Siphonina, 100
 jacalitosana, Dosinia ponderosa, 104
 japonica, Mya (Mya) arenaria, 105
 jenkinsi, Odontopleura, 119
 jenkinsi, Scutellum, 120
 jewettii, Hyalina (Cystiscus), 112
 jewettii, Turritella, 116
 jordani, Neptunea (Colus), 114
 jordani, Pecten (Pecten) islandicus, 107
 jordani, Pteria, 108
 juanensis, Chione, 103
 juanensis, Conus, 111
 kelletii, Kelletia (Kelletia), 112
 kernensis, Pholadomya, 108
 kewi, Scutaster vaquerosensis, 123
 labiata, Textularia, 100
 lakei, Arca, 102
 lamellosa, Cumingia, 103
 latejugata, Nodosaria, 100
 latiauratus, Pecten (Aequipecten), 106
 lamellifer, Irus, 104
 latrania, Pinna, 108
 lepisima, Acmaea, 110
 lewisi, Gyreneum, 112
 lohri, Pecten (Patinopecten) healeyi, 107
 loismartinae, Cylichna, 112
 longidens, Dosinia ponderosa, 104
 longior, Anatina (Raeta) plicatella, 101
 longispina, Ceratocephala, 118
 longispinifex, Scutellum, 120
 lovenioides, Megapetalus, 122
 lucida, Siliqua, 108
 lugubris, Morula (Morunella), 113
 lurida, Tritonalia, 116
 luteola, Corbula (Lentidium), 103
 lyra, Scissurella, 115

macropleurus, Loganellus, 119
 macroschisma, Pododesmus, 108
 magister, Forreria, 112
 major, Phillipsia, 119
 mamillaris, Calyptarea, 111
 margaritana, Dosinia, 103, 104
 mariposa, Turbonilla (Pyrgolampros), 116
 marquerensis, Pecten (Patinopecten), 107
 mendicus, Nassarius (*Schizopyga*), 113
 mendicus, Nassarius (*Tritia*), 114
 mercedensis, Moniliopsis graciosa, 113
 merriami, Drillia, 112
 merriami, Solenosteira, 115
 mexicana, Eponides, 100
 mexicanus, Robulus, 100
 microsperma, Nucula (Ennucula), 105
 milowi, Cytherelloidea, 121
 minuscula, Glans, 104
 minuta, Xestoleberis, 122
 modesta, Admete, 110
 moesta, Macoma, 104
 monotimeris, Pecten (Aequipecten) latiauratus, 106
 montereyana, Turritella, 117
 multicostata, Arca (Arca), 102
 munda, Tritonalia lurida, 116
 nasuta, Macoma, 104
 nipponica, Thyasira bisecta, 109
 nobilis, Architectonica, 110
 nosoniensis, Griffithides, 119
 nudicostatus, Robulus mexicanus, 100
 nuttalli, Saxidomus, 108
 nuttallii, Lucina (Myrtea), 104
 nuttallii, Schizothaerus, 108
 obesa, Cancellaria, 111
 obstipa, Balcis (Vitreolina), 110
 ocoyana, Ficus (Trophosycon), 112
 ocoyana, Turritella, 117
 octocamerata, Gyroidina soldanii, 100
 oena, Platycosta, 122
 oldroydi, Tellina, 109
 olivaceous, Melampus, 113
 onyx, Crepidula, 112
 orcutti, Sanguinolaria (Nuttallia), 108
 oregonensis, Ranella (Priene), 115
 ornata, Griffithides, 119
 orpheus, Trophon (Boreotrophon), 116
 orthomorpha, Mactra (Mactra), 105
 oweniana, Conus, 111
 painei, Acteon (Rictaxis), 110
 pallidus, Clavus (Cymatosyrinx), 111
 panzana, Macoma, 104
 panzana, Oculina, 101
 panzana, Pedalion, 108
 partschi, Scutellum, 120
 parva, Lloydia, 119
 parviusculus, Proetus, 120
 patula, Siliqua, 108
 pedroana, Olivella, 114
 pedroana, Spirotropis (Antiplanes) perversa, 115
 penicillata, Pseudomelatoma, 114
 penita, Pholadidea (Pholadidea), 108
 percarus, Pecten (Aequipecten), 106
 perdisparis, Arca, 102
 perpinguis, Nassarius (*Schizopyga*), 114
 perpinguis, Nassarius (*Tritia*), 114
 perplexa, Hesperocidaris, 122

perrini, Spondylus, 109
 pertumida, Turritella inezana, 116
 peruviana, Anomia, 101
 perversa, Spirotropis (Antiplanes), 115
 phanea, Odostomia (Evalea), 114
 planolata, Eucytherura, 121
 planospira, Cancellaria obesa, 111
 pleistocenea, Randallia, 122
 pleurexanthemus, Ceraurus, 118
 plicatella, Anatina (Raeta), 101
 poulsoni, Tritonalia, 116
 ponderosa, Dosinia, 104
 prelamellifer, Iirus lamellifer, 104
 procumbens, Arca, 102
 projecta, Dosinia (Dosinidia) margaritana, 103
 pseudotrigona, Paijenborchella, 122
 psila, Discocyclina, 100
 pufferi, Vasum (Vasum), 117
 punctata, Pandora, 106
 punctocostata, Puncturella, 114
 punctulum, Mistostigma, 113
 purissima, Pecten (Patinopecten), 107
 purpuratus, Pecten (Aequipecten), 106
 pusillanima, Turbinolia, 101
 putnami, Oligopygus, 123
 quinquecincta, Moniliopsis incisa, 113
 ralphi, Puncturella, 115
 rana, Phacops, 119
 rattei, Odontopleura, 119
 rattei, Proetus, 120
 recurva, Strombina, 115
 remondii, Surculites (Megasurcula), 115
 renaudi, Spirotropis (Typhlomangelia), 115
 restorationensis, Paphia, 106
 restorationensis, Venerupis (Protorthaca), 109
 rex, Galeodea, 112
 rivulata, Arca, 102
 rogersi, Sphaerium (Amesoda), 108
 rostrata, Calymene, 118
 rostrata, Calymenella, 118
 rotundomontana, Chrysallida, 111
 ruderata, Venerupis (Protorthaca) staminea, 109
 rugatum, Bittium (Semibittium), 110
 ruginodosa, Ficus (Trophosycon) ocoyana, 112
 sampsoni, Phillipsia, 119
 sandiegensis, Cibicides, 99
 sandiegensis, Gaudryina convexa, 100
 santana, Arca (Andara), 102
 santana, Olivella, 114
 santana, Terebra, 116
 santaclarana, Arca (Andara), 102
 santacruzana, Turcicula, 116
 schenki, Cardium, 103
 scutulata, Littorina, 113
 sechurana, Chione, 103
 secta, Macoma, 104
 securis, Venus (Chione), 109
 semiinflata, Pseudomelatoma penicillata, 114
 semirostrata, Nucula (Acila), 105
 septentrionalis, Glycymeris, 104
 serrai, Rapana, 115
 serratus, Phacops, 119
 sespeensis, Arca, 102
 sespeensis, Macoma, 105
 sespeensis, Turritella inezana, 116
 shumanensis, Thais (Nucella), 116

- sicarius, Solen, 108
 simplicius, Cerithium, 111
 slevini, Pecten (Janira) bellus, 107
 soldanii, Gyroidina, 100
 solidula, Lacuna divaricata, 112
 spadicea, Cypraea, 112
 spinata, Eucytherura, 121
 spirata, Acanthina, 109
 squamulifer, Ocenebra, 114
 squamulifer, T.[rophon], 116
 stalderi, Amiantis callosa, 101
 staminea, Venetupis (Protothaca), 109
 staminea, Venus (Protothaca), 109
 stantoni, Macrocallista, 105
 stearnsii, Pecten (Janira) stearnsii, 107
 stephensi, Atrina, 102
 stephensi, Callianassa, 121
 stevensoni, Phillipsia, 119
 stocki, Nassarius, 114
 strongi, Arca (Barbatia), 102
 strongi, Gyreneum (Bechtelia), 112
 stultorum, Tivela, 109
 subacuminata, Bulimina, 99
 subcalva, Bulimina, 99
 subdiaphana, Clementia (Compsomyax), 103
 subdolus, Pecten (Aequipecten) purpuratus, 106
 succincta, Venus (Chione), 109
 sulcata, Turbinolia, 101
 swiftii, Pecten (Pallium), 107
 taphria, Nuculana, 105
 tapinus, Spatangus, 123
 tayloriana, Ostrea, 105
 temblörensis, Cylichna, 112
 temblörensis, Turritella, 117
 tenuis, Encope, 122
 tenuis, Panope, 106
 tenuisculpta, Lucina (Myrtea), 104
 terminopunctata, Cytherella, 121
 terminumbonis, Arca (Navicula), 102
 terfa, Balcis (Balcis), 110
 trapezoides, Thracia (Thracia), 109
 triangularis, Quinqueloculina, 100
 tridesmia, Clathurella (Glyphostoma), 111
 triebeli, Idiocythere, 121
 tritonidea, Cancellaria, 111
 tuberosa, Mitrella, 113
 turgidulum, Eocytheropteron, 121
 umbilicatus, Nonion, 100
 unisulcata, Richardsonia, 120
 valentinei, Chione, 103
 vancouverensis, Arca, 102
 vancouverensis, Parallelodon (Nanonavis), 106
 vancouverensis, Pecten (Pseudamusium), 108
 vancouverensis, Laqueus, 101
 vanveleki, Turritella, 117
 vanwinkleae, Pecten, 108
 vaquerosensis, Anomia, 102
 vaquerosensis, Antigona, 102
 vaquerosensis, Ostrea, 106
 vaquerosensis, Scutaster, 123
 variata, Turritella, 117
 varicostatum, Epitonium (Opalia), 112
 ventricosa, Cardita, 103
 venturana, Ostrea, 106
 verrucosa, Semele, 108
 versabilis, Eucytherura, 121
 versicolor, Amphissa, 110
 vespertina, Ostrea, 106
 vigilans, Vogdesia, 120
 vizcainoensis, Dendraster, 122
 vogdesi, Bumastus, 117
 vogdesi, Calymene, 118
 washingtonianus, Viviparus, 117
 weaveri, Miopleiona, 113
 whitei, Ferminalscala, 112
 whitneyi, Nassarius (Uzita) arnoldi, 114
 wilkesanus, Chicoreus (Murithais), 111
 xantusi, Lucina (Miltha), 104
 yassensis, Cyphaspis, 118
 ynezana, Ostrea eldridgei, 105
 ynezana, Tritonalia, 116
 ynezanus, Conus oweniana, 111

